

Review

UAVs as remote sensing platforms in plant ecology: review of applications and challenges

Zhongyu Sun¹, Xiaonian Wang², Zhihui Wang¹, Long Yang¹, Yichun Xie^{3,*}  and Yuhui Huang⁴

¹Guangdong Open Laboratory of Geospatial Information Technology and Application, Guangzhou Institute of Geography, Guangdong Academy of Sciences, Guangzhou 510070, China, ²School of Geographical Sciences, South China Normal University, Guangzhou 510631, China, ³Department of Geography and Geology, Eastern Michigan University, Ypsilanti, MI 48197, USA, ⁴Guangdong Provincial Key Laboratory of Silviculture, Protection and Utilization, Guangdong Academy of Forestry, Guangzhou 510520, China

*Corresponding author. E-mail: yxie@emich.edu

Handling Editor: Shaopeng Wang

Received: 26 March 2021, **First Decision:** 30 June 2021, **Accepted:** 29 July 2021, **Online Publication:** 20 August 2021

Abstract

Aims Unmanned aerial vehicles (UAVs), i.e. drones, have recently emerged as cost-effective and flexible tools for acquiring remote sensing data with fine spatial and temporal resolution. It provides a new method and opportunity for plant ecologists to study issues from individual to regional scales. However, as a new method, UAVs remote sensing applications in plant ecology are still challenged. The needs of plant ecology research and the application development of UAVs remote sensing should be better integrated.

Methods This report provides a comprehensive review of UAV-based remote sensing applications in plant ecology to synthesize prospects of applying drones to advance plant ecology research.

Important Findings Of the 400 references, 59% were published in remote sensing journals rather than in plant ecology journals, reflecting a substantial gap between the interests of remote sensing experts and plant ecologists. Most of the studies focused on UAV remote sensing's technical aspects, such as data processing and remote sensing inversion, with little attention on answering ecological questions. There were 61% of studies involved community-scale research. RGB and multispectral cameras were the most used sensors (75%). More ecologically meaningful parameters can be extracted from UAV data to better understand the canopy surface irregularity and community heterogeneity, identify geometrical characteristics of canopy gaps and construct canopy chemical assemblies from living vegetation volumes. More cooperation between plant ecologists and remote sensing experts is needed to promote UAV remote sensing in advancing plant ecology research.

Keywords UAVs, drones, unmanned aircraft systems (UASs), plant ecology, species identification, community function


无人机遥感在植物生态学中的应用与挑战

摘要: 无人机为获取高时空分辨率的遥感数据提供了经济灵活的工具, 为植物生态学家开展从个体到区域尺度的生态学研究提供了新的机遇和手段。但作为一种新兴的技术手段, 当前无人机遥感在植物生态学中的应用仍充满了挑战, 植物生态学的科研需求与无人机遥感的生态应用需要更为深入的融合。

本文综述了无人机遥感技术在植物生态学中的应用，展望了无人机在植物生态学研究中的应用前景。在所综述的400篇文献中，59%的文章发表于非植物生态学领域的遥感类期刊，遥感学者与生态学者的关注点存在较大差异。当前的研究集中在无人机遥感的技术层面，如数据处理和遥感反演方法等，对生态学问题本身的关注较少。综述的文献中，61%的研究案例集中在群落尺度，可见光(RGB)相机和多光谱相机是最常用的传感器类型(75%)。无人机遥感数据中蕴藏着诸多待挖掘的、有重要意义的生态参数，这些参数有助于我们识别林窗的几何特征，构建林冠的化学组合，更好地了解林冠表面的不规则性和群落的异质性。无人机遥感技术在植物生态学研究中的深入应用，需要集植物生态学家和遥感专家之合力共同推进。

关键词：无人机，植物生态学，物种鉴定，尺度，功能性状

Graphical Abstract



INTRODUCTION

Plant ecologists study the relationships between plants and their environment from gene to global scales (Keddy 2007). Remote sensing from spaceborne, airborne and terrestrial platforms has provided abundant data and analytical tools for plant ecology studies at regional and global scales (Myneni and Ross 2012; Xie *et al.* 2008). As early in 2018, 1738 satellites in Earth orbit were equipped with various types of remote sensors, such as multispectral cameras, hyperspectral cameras and light detection and ranging (LiDAR) sensors (Union of Concerned Scientists 2018). The satellites collect data with spatial resolutions ranging from kilometers (e.g. NOAA-AVHRR, 1100

m) to submeters (e.g. Worldview III, 0.31 m). The applications include the characterization of vegetation type, aboveground biomass, leaf area index (LAI), vegetation cover and canopy chemistry at regional and global scales (Gomez *et al.* 2019). Although these data have enabled researchers to assess ecological conditions in the context of global environmental change, satellite and airborne-based remote sensing systems often fail to meet the requirements of ecological and environmental research. Use of these systems by plant ecologists is limited by inadequate spatial, temporal and spectral resolution; lack of operational flexibility and noise caused by atmospheric conditions (Adam *et al.* 2010; Huylenbroeck *et al.* 2020). For ground-based remote sensing, with handheld devices or

tower crane equipment, its measurements are often taken at limited points and the measurement process is time consuming. Accordingly, it is hard to apply the ground-based remote sensing to complex environments over larger areas.

In recent years, unmanned aerial vehicles (UAVs), also known as remotely piloted aircraft system, unmanned aircraft systems or drones, have been widely used in plant ecology. Such UAVs are easy to deploy and are economical and most importantly, technically capable of collecting imagery data at fine spatial, spectral and temporal resolutions. UAV data can complement ground observations and data collected from aircraft and satellite remote sensing platforms, and thereby provide a comprehensive remote sensing system for plant ecology studies ranging from individuals to ecosystems (Singh and Frazier 2018; Torresan *et al.* 2017; Valbuena *et al.* 2020).

There is an increasing number of studies on the applications of UAV remote sensing, and these studies span a broad array of topics including UAV platform classification and development (Colomina and Molina 2014; Floreano and Wood 2015; Hassanalain and Abdelkefi 2017; Watts *et al.* 2012), UAV applications in agriculture (Perich *et al.* 2020; Yang *et al.* 2018; Zhang and Kovacs 2012), resource management (Oliveira *et al.* 2020; Shahbazi *et al.* 2014), environmental studies (Pichon *et al.* 2019; Wang *et al.* 2018; Whitehead and Hugenholtz 2014; Whitehead *et al.* 2014) and biodiversity monitoring (Bagaram *et al.* 2018; Guo *et al.* 2016b). Some pioneering studies also discussed the potential use of UAV remote sensing in ecology (Anderson and Gaston 2013; Lian and Wich 2012; Valbuena *et al.* 2020). In this report, we review UAV remote sensing systems and their applications in plant ecology from a perspective that integrates the views of ecologists and remote sensing professionals. Our analyses are divided into five levels, i.e. individuals, populations, communities, ecosystems and landscape. We conclude our review by discussing the challenges and prospects of UAV remote sensing in plant ecology research.

REVIEW METHODS

We collected data from the ISI Web of Science Core Collection using the following search terms: TS = ('remotely piloted aircraft system') or ('unmanned aerial vehicles') or TS = ('unmanned aircraft systems') or TS = (drones) or TS = ('unmanned aerial systems') and LANGUAGE: (English). The

search results were then refined according to WEB OF SCIENCE CATEGORIES: (Remote Sensing or Ecology or Agriculture Multidisciplinary or Plant Sciences or Forestry). TIME SPAN was set at 2004–20 (August 2020), and INDEXES was set at SCI-EXPANDED. With the article type limited to 'research articles' and 'review articles', we found 1425 records in the ISI Web of Science. We screened the abstracts of references and removed those records that were not relevant to this review. In total, we identified 400 papers for detailed review, which included 354 research articles and 46 review papers (Appendix S1: Reviewed references list). Finally, our database included 354 reports of original research.

We designed a standardized template to review these articles (Appendix S1: Template of reviewed studies). The template included the following criteria: published year, the institution of the first author, location of the study site(s), study area, vegetation form, observation scale, UAV type, UAV producer, sensor type, flying altitude, processing method and research objectives in the context of plant ecology.

The review is presented in four parts. The first part is titled 'UAV Systems, UAV Data Processing and Analysis', which provides basic information about the UAV instrumentation, the data and the related processing and analytical methods. The second part is called, 'Applications of UAV Remote Sensing in Plant Ecology'. This part focuses on the relevance of UAV technology to ecology through broad applications of UAV remote sensing in plant ecology. This part is organized according to well-acknowledged contributions of UAV to plant ecological studies and is also structured by considering three primary types of plant ecological studies: (i) Individual to population scales: individual detection, physiological assessment and species classification and distribution; (ii) Community scale: composition, structure, foliar functional traits, biodiversity and biomass and (iii) From ecosystem to landscape scale: monitoring and management. The third part discusses the challenges and prospects of UAV remote sensing in plant ecology. The main technical challenge is how to effectively fuse multisource remote sensed data including UAVs in the context of supporting plant ecology studies, while the primary application challenge is how to better integrate UAV obtained data to answer or solve basic scientific questions facing plant ecology. The prospects of UAV remote sensing in plant ecology are very promising, including automatic species identification, multiscaling spatial exploration of plant ecosystems, increased UAV applications from

describing ecological phenomenon to answering ecological questions, and the needs of more UAV-based novel methods to answer ecological questions. The last is the section of Conclusions.

UAV SYSTEMS, UAV DATA PROCESSING AND ANALYSIS

UAV systems

UAV remote sensing systems have at least five components, i.e. a platform system, a sensor system, a ground control and data transmission system, a data processing system and operators (Fig. 1). Previous reviews of UAV platforms and sensing payloads can be found in Watts *et al.* (2012) and Hassanalian and Abdelkefi (2017). In Table 1, we have summarized the advantages of UAV remote sensing systems by comparing them with traditional spaceborne and airborne remote sensing systems (Table 1). The first advantage is low cost (Xie *et al.* 2015; see a comparable cost analysis in Appendix S2). As a second advantage, UAV remote sensing provides high temporal and spatial resolutions. UAV systems

can acquire imagery with centimeter resolution at almost any time of the day and under most weather conditions. The very high-resolution imagery makes it possible for ecologists to study many canopy properties, including canopy structure and dynamics. Another advantage is that the operation of a UAV remote sensing system is flexible. In a complicated environment, small UAVs can take off and land on an operator's hands, which greatly increases the utility of UAV remote sensing in ecological studies. In addition, UAV remote sensing systems are relatively easy to use, such that operation requires only a short training period. As a final advantage, the ultra-low altitude flying of UAVs can reduce the effect of cloud on imagery and thereby improve the data quality.

Most of the 354 case studies of UAV application in plant ecology used a rotary wing (64%) or a fixed-wing UAV platform (26%). About 88% of these were off-the-shelf UAV platforms such as the DJI phantom series and the senseFly eBee series (Fig. 2a and b). Some parafoil wing and vertical take-off and landing fixed-wing UAVs were used, but the percentages were very low. In the 354 studies, the UAVs usually flew at altitudes of 10–120 m in order to obtain images with




Figure 1: The components of a UAV remote sensing system (adapted from Sun *et al.* 2017).

Table 1: Characteristics of spaceborne, airborne and UAV remote sensing systems

System	Spatial resolution	Swath	Controllability	Load	Flight time	Cost	Flexibility	Degree of cloud influence
Space remote sensing	1–25 m	10–50 km	—	—	—	Very high	Lowest	Moderate
Aerial remote sensing	0.1–2 m	0.5–5 km	Requires a pilot and a high-quality landing point	No limitation	Long	High	Low	Low
UAV remote sensing	0.5–10 cm	50–500 m	Remote control or fly automatically, nearly no demand for a landing point	With limitation	Short	Low	Highest	Very low

ultra-high resolution and comply with UAV aviation regulations (Fig. 2c).

UAV data processing and analysis

Various types of sensors are available for the UAV-based platform, including RGB, multispectral, hyperspectral, thermal and LiDAR sensors. RGB cameras are most commonly used due to their low costs, lightweights and ease of use (Bagaram *et al.* 2018; Cunliffe *et al.* 2016; Pichon *et al.* 2019). Multispectral sensors (e.g. MicaSense RedEdge 3 camera, Micasense, WA, USA) provide more spectral bands (e.g. red-edge: 760 nm; near-infrared [NIR]: 810 nm) which can better evaluate plant health and stress status (Adam *et al.* 2010; Baluja *et al.* 2012; Wang *et al.* 2019b). Hyperspectral sensors (e.g. Cubert S185, Cubert, Ulm, Germany) provide continuous spectrum with narrow bandwidths (<10 nm) and thus offer useful means of detecting fine absorption features of plant biochemicals (Kwon *et al.* 2020; Pölonen *et al.* 2013; Saarinen *et al.* 2018). Thermal infrared sensors capture the thermal radiation from the plants which can be used to estimate the surface temperature of plants for monitoring plant water stress and forest fire (Calderón *et al.* 2013; Messina and Modica 2020; Smigaj *et al.* 2017). LiDAR sensors measure the distance to a target by travel time of the emitting laser light and can characterize canopy structural parameters such as tree height, crown width and canopy cover (de Almeida *et al.* 2020; Ganz *et al.* 2019).

In general, the data preprocessing of optical sensors (RGB, multispectral, hyperspectral and thermal sensors) includes geometric correction and radiometric correction. Geometric correction is generally performed based on the GPS and inertial measurement unit data. Ground control points are often used to improve the accuracy of geometric corrections. Recently, new photogrammetry techniques such as structure from motion (SfM) have been used to generate orthophoto mosaic based on matching feature across overlapped images (Gil-Docampo *et al.* 2020; Wallace *et al.* 2016). Software such as Agisoft Photoscan (now Metashape, Agisoft LLC, St. Petersburg, Russia) and Pix4d mapper (Pix4D SA, Lausanne, Switzerland) have integrated these algorithms into software and support automatic image preprocessing, which include image matching, mosaic, geometric correction, brightness and contrast adjustment (Formoo *et al.* 2019).

Radiometric correction converts the digital numbers of images to reflectance. RGB and




Figure 2: Characteristics of UAVs used in plant ecology studies (**a** and **b**), including their flying altitudes (**c**).

thermal cameras are generally hard to calibrate due to the nonlinear gamma correction effect. As a result, they are often used for image classification or temperature measurement which does not highly rely on radiometric calibration. Multiple spectral sensors have different ways to perform radiometric correction. One way is to have a sensor onboard to measure the downwelling radiance and calculate in-time reflectance. Another way is to place one or more ground targets with standard known reflectance during the flight and then to perform an empirical line correction (Aasen *et al.* 2018). It should be mentioned that the empirical line correction method requires a stable sunny

sky condition, i.e., the solar illumination does not change much during the flight. For long distance flights, a concurrent radiance measurement of ground targets using field spectrometers is often suggested (Aasen *et al.* 2018). These techniques of radiometric corrections also apply to hyperspectral sensors.

Data processing and analysis include imagery-based methods (for RGB data), point clouds methods (for RGB and LiDAR data), statistical models (for multispectral and hyperspectral data) and physical models (for multispectral and hyperspectral data). RGB images can be used to detect trees using object-based image analysis such as eCognition software or deep learning methods (Mu *et al.* 2018; Petrich *et al.* 2020). The RGB images can also be used to observe forest phenology and crop lodging (Berra *et al.* 2019; Zhang *et al.* 2020).

Individual tree segmentation can be achieved from LiDAR and RGB mosaic data (Wallace *et al.* 2016). LiDAR and RGB point clouds are generated in different ways. LiDAR point clouds can be obtained during the laser scanning, but RGB point clouds need to be constructed through the SfM algorithm. After the point cloud generation, LiDAR and RGB data can be further processed using the same methods. The point clouds are first interpolated to create digital surface model (DSM) and digital terrain model (DTM) by using techniques such as Delaunay Triangulation and triangulated irregular networks. The height calculation is done with the canopy height model (CHM) by subtracting DTM from DSM. Tree detection can then be conducted from CHM or DSM using variable-sized window and watershed delineation (Yin and Wang 2019), or directly from point clouds using point cloud segmentation and layer stacking (Wallace *et al.* 2016). Tree height, crown width and canopy cover can be derived from CHM or point clouds (Guerra-Hernandez *et al.* 2016; Solvin *et al.* 2020). RGB data usually require more interpolations to build DSM and DTM, because the point clouds obtained through the SfM algorithm are much sparser than those from LiDAR data. Therefore, the structural attributes extracted from the LiDAR point clouds are usually more accurate than those extracted from the RGB point clouds.

Statistical models are developed for classification and regression. Machine learning algorithms such as random forest, support vector machine and convolutional neural networks have been developed to perform automatic (unsupervised) or

semiautomatic (supervised) classification of UAV remote sensing data (Nguyen *et al.* 2019). Recently, deep learning algorithms based on neural network have emerged as an effective tool for species classification (Lopez-Jimenez *et al.* 2019; Plesoianu *et al.* 2020; Zou *et al.* 2019). For regression, the relationship between a parameter of interest (e.g. leaf chlorophyll or nitrogen) and the spectral data is established. Statistical approaches include vegetation indices, linear regression approaches such as stepwise linear regression and partial least squares regression, nonlinear regression approaches such as random forest, support vector machine and artificial neural network (Padua *et al.* 2017). Physical models describe the interaction and transfer of solar radiation based on physical laws and provide the advantage of transferability over statistical models. Physical model was rarely used in previous studies and was found to map the reflectance anisotropy of a potato canopy (Roosjen *et al.* 2017).

APPLICATIONS OF UAV REMOTE SENSING IN PLANT ECOLOGY

General characteristics of UAV remote sensing applications in plant ecology

The 354 case studies in our database were carried out in 43 countries, mostly in North America, East Asia and West Europe; the two leading countries were the USA and China (Fig. 3). The earliest study in our database was carried out in 2005 in the USA; after that, the number of studies increased rapidly over time, especially during 2016–2019 (Appendix S2; Supplementary Fig. S1). There were 215 case studies published in remote sensing journals, while 139 in plant science and ecology journals.

Among the 46 review papers, 21 were published in remote sensing journals and 25 were published in plant science and ecology journals. The reviews in remote sensing journals mainly focused on the challenges and applications of UAV systems in environmental monitoring. Reviews in plant science and ecology journals primarily focused on precision agriculture and forest management. The previous reviews were conducted from the perspective of remote sensing experts, i.e. they focused on mapping, measuring and monitoring plant properties rather than on answering questions in biology or ecology. There is an obvious gap between the concerns of remote sensing scientists and ecologists. Of the 354 case studies, most of them used RGB or

multispectral cameras, focused on the community to ecosystem scale and applied on forests and crops (see a detailed tabulation in Appendix S1). Therefore, a comprehensive assessment of UAV applications in plant ecology is needed, especially for different ecological scales such as the individual, population, community, ecosystem and landscape scale.

Individual to population scales: individual detection, physiological assessment and species classification and distribution

UAV remote sensing data have been widely used to measure crown characteristics of individual plants. Surovy *et al.* (2018) built a point cloud to estimate the height and position of individual trees. Mu *et al.* (2018) measured the crown width and crown projection area of individual peach trees. Stress and other physiological changes in plants can be tracked by variations in the visible and NIR wavelengths. Stressed plants often exhibit a higher reflectance in the visible and lower reflectance in the NIR than nonstressed plants. Indicators of plant stress and growth such as leaf chlorophyll and LAI can also be estimated from the spectra. When plants experience drought stress, the leaf stomata close, resulting in a higher leaf temperature, which can be monitored using thermal imagery (Leinonen *et al.* 2006). For instance, Berni *et al.* (2009) used thermal imagery to assess the water-stressed status of peach trees and to guide precision irrigation (Berni *et al.* 2009). Spectral indices, such as the normalized difference vegetation index ($NDVI = (R_{800} - R_{670}) / (R_{800} + R_{670})$, Rouse *et al.* 1974), the renormalized different vegetation index ($RDVI = (R_{800} - R_{670}) / \sqrt{(R_{800} + R_{670})}$, Rougean and Breon 1995), the modified triangular vegetation index 1




Figure 3: The percent of case studies in our database conducted in the indicated countries.

($MTVII = 1.2 \times [1.2 \times (R_{800} - R_{550}) - 2.5 \times (R_{670} - R_{550})]$), Haboudane *et al.* 2004), the triangular vegetation index ($TVI = 0.5 \times [120 \times (R_{750} - R_{550}) - 200 \times (R_{670} - R_{550})]$), Haboudane *et al.* 2004), the ratio of transformed chlorophyll absorption ratio index ($TCARI = 3 \times [(R_{700} - R_{670}) - 0.2 \times (R_{700} - R_{550}) \times (R_{700}/R_{670})]$), Haboudane *et al.* 2002) and the optimized soil-adjusted vegetation index ($OSAVI = (1 + 0.16) \times (R_{800} - R_{670}) / (R_{800} + R_{670} + 0.16)$), Haboudane *et al.* 2002), have been used to assess the water stress of orange and mandarin trees (Zarco-Tejada *et al.* 2012) and grapevines (Baluja *et al.* 2012; Romero *et al.* 2018). Hyperspectral and thermal imageries have also been used to quantify the effect of a disease (Verticillium wilt) on the stomatal conductance of olive leaves (Calderón *et al.* 2013). Species classification is the most common use of UAV remote sensing at the individual scale. Satellite remote sensing has been widely used to generate global or national land cover maps on which the vegetation is classified as forests, grasslands, deserts and wetlands (He *et al.* 2009). The image processing and classification procedures have been detailed by Laliberte *et al.* (2011). Because of the coarse spatial resolution of satellite imagery, however, plant species are difficult to be distinguished unless the target species have unique growth forms or phenology (Bradley 2014; Huang and Asner 2009). In contrast, UAVs can obtain ultra-high spatial and spectral resolution imagery, satisfying the requirements of vegetation classification at the species level. In a lake ecosystem, e.g., 49 lacustrine plant species/vegetation classes were identified by using high-resolution optical imagery with an accuracy of 95.1% (Husson *et al.* 2013). By incorporating tree heights with spectral features, image textural features and hyperspectral vegetation indicators, UAV accuracy in identifying mangrove species can be as high as 88% (Cao *et al.* 2018). Understorey herbs can potentially be identified to species in sparse forests with optical and multispectral imagery (Leduc and Knudby 2018; Sanders 2017).

The accuracy of species identification depends on four factors: spatial resolution, spectral resolution, habitat complexity and classification algorithms. Spatial resolution is the primary determinant of identification accuracy (Ashraf *et al.* 2010). A 5-cm resolution was not sufficient to identify some herbs at the species level (Dunford *et al.* 2009). In contrast, images with 1-cm resolution could identify herbaceous plant species in a wetland with an accuracy of 93% (Ishihama *et al.* 2012). At a scale

of 1/50 unit, populations of *Phragmites australis*, *Typha domingensis* and *Miscanthus sacchariflorus* were clearly discriminated in a river estuary (Kaneko and Nohara 2014). Therefore, ultra-high resolution images obtained with cameras on UAVs flying at a low altitude will greatly improve plant identification at the species level (Cao *et al.* 2018; Li *et al.* 2017; Yang *et al.* 2016; Zarco-Tejada *et al.* 2012). However, the high cost of hyperspectral UAVs (Manfreda *et al.* 2018; Whitehead and Hugenholtz 2014) limits their applications in plant ecology (Adão *et al.* 2017).

Accurately identifying species in complicated habitats is challenging, especially when the target species are small and similar to each other. A case study in the Arctic Tundra indicated that VIS–NIR high-resolution imagery could identify the main vegetation groups but could not distinguish between species (Mora *et al.* 2015). In rain forests and subtropical forests, tree properties (e.g. crown size, crown status, crown contour, crown architecture, foliage cover, foliage texture and foliage color) must be integrated in order to identify the dominant species in the canopy layer, but the number of species that can be accurately identified is still limited (Trichon 2001; Yang *et al.* 2016). In summary, UAV remote sensing for species identification has mostly been applied in relatively simple habitats, such as rangelands (Karl *et al.* 2020; Laliberte *et al.* 2011; Rango *et al.* 2006, 2009), wetlands (Chabot and Bird 2013; Doughty and Cavanaugh 2019; Husson *et al.* 2014; Ishihama *et al.* 2012; Zweig *et al.* 2015), plateau shrub swamps (Fletcher and Erskine 2012) and riparian forests (Husson *et al.* 2013). There are a few studies carried out in urban areas although the habitats are relatively simple. Flying drones in cities often requires strict approval procedures, which limits their use in urban environments.

Image processing methods are required for the use of UAV remote sensing in plant ecology. In addition to popular pixel-based and object-based image analysis, a ‘feature learning’ approach based on machine learning represents a novel and effective method for species identification (Hung *et al.* 2014; Lary *et al.* 2016; Plesoianu *et al.* 2020). The fusion of multisource data, such as hyperspectral, RGB and LiDAR data, can also improve the accuracy of species classification (Cao *et al.* 2018; Yin and Wang 2019).

Mapping the spatial distribution of plant species at the population scale is a popular application of UAV remote sensing. For instance, Flynn and Chapra (2014) used UAV remote sensing to map the

distribution of a green alga (*Cladophora glomerata*) in rivers. Kalacska *et al.* (2013) used UAV remote sensing to map the spatial distribution of *Eriophorum vaginatum* and to evaluate its contribution to CH₄-C flux in an ombrotrophic bog in Canada. Other researchers have used UAV remote sensing to detect and map species and thereby to help manage populations of weeds and invasive plants (Abeyasinghe *et al.* 2019; Alvarez-Taboada *et al.* 2017; Hill *et al.* 2017; Peña *et al.* 2013; Tamouridou *et al.* 2017).

Community scale: composition, structure, foliar functional traits, biodiversity and biomass

Species composition (13 species) of a wetland area in Hong Kong was mapped with a UAV-based hyperspectral image and DSM derived from photogrammetric point clouds (Li *et al.* 2017). Banerjee *et al.* (2017) identified and mapped five plant species in a complex upland swamp community using a UAV-hyperspectral system, and the overall accuracy of classification was 88.9%. In a study by Chisholmryan *et al.* (2013), the use of a UAV LiDAR system (flown 1.5 m above the ground) to obtain DBH data for trees provided a new way to conduct below-canopy forest surveys.

The horizontal as well as the vertical structure of plant communities can be investigated through UAV remote sensing (Campos-Vargas *et al.* 2020; Schneider *et al.* 2019). Using a UAV-optical camera, Getzin *et al.* (2012) characterized the horizontal patterns of small gaps (<5 m²) in 10 temperate forests and found that these small gaps, which could hardly be detected by conventional aerial or satellite images, made up the majority of gaps in the canopies. The canopy height, aboveground biomass and canopy complexity measured by UAV LiDAR and an optical camera were used to estimate frugivorous bird abundance and forest recovery (Zahawi *et al.* 2015). In areas with relatively low canopy closure, the SfM point clouds obtained with a UAV-optical camera provided abundant information on forest structure (Jensen and Mathews 2016; Wallace *et al.* 2016). LAI and vegetation coverage can be estimated with UAV remote sensing to evaluate community structure (Tian *et al.* 2017; Wang *et al.* 2019a). UAV remote sensing has also become increasingly important in forest phenology studies (Mariano *et al.* 2016). The spatial and temporal changes of a forest were studied at the community scale by using a UAV-optical camera (Klosterman *et al.* 2018).

Foliar functional traits refer to a range of biochemical and physiognomic characteristics of

plants, such as macronutrients (N, P, K, Ca, Mg, S), trace minerals (B, Cu, Fe, Mn, Zn) and Al, cellulose, lignin, sugars and starches, have been widely estimated using manned aircraft remote sensing (Asner *et al.* 2014; Schneider *et al.* 2017; Wang *et al.* 2018, 2020). With higher security and flexibility, UAV remote sensing has the potential to collect foliar functional traits with higher spatial and temporal resolutions. A number of studies have been conducted in croplands and grasslands (Table 2).

The diversity of the dominant species in the upper canopy layer of a subtropical or mangrove forest were quantified using UAV-optical or hyperspectral cameras (Cao *et al.* 2018; Yang *et al.* 2016). Theoretically, the biodiversity of a canopy layer can be easily calculated based on the information of tree canopies. However, UAV remote sensing has rarely been used to directly measure biodiversity (Guo *et al.* 2016b). Saarinen *et al.* (2018) used UAV-based photogrammetric point clouds and hyperspectral imaging to monitor dead wood quantity and species richness in a boreal forest. Getzin *et al.* (2012) used UAV to calculate eight different gap metrics and to determine whether those metrics were correlated with floristic biodiversity of the forest understory at a landscape scale.

Plant biomass can be estimated through UAV remote sensing in two ways (Man *et al.* 2014). For the first approach, a relationship between the remote sensing data and biomass is first established; biomass can then be estimated by *K* nearest neighbor classification, multiple regression analysis, neural network methods or statistical ensemble methods. For example, vegetation indices derived from UAV-multispectral or hyperspectral images have been widely used to estimate aboveground biomass, productivity or yield (Geipel *et al.* 2014; Getzin *et al.* 2012; Gonzalez-Jaramillo *et al.* 2019; Pölonen *et al.* 2013). On the other hand, tree height, DBH or crown volume are extracted from UAV remote sensing images and then used to calculate biomass with allometric equations (Bendig *et al.* 2014; Cunliffe *et al.* 2016). In general, UAV LiDAR can generate more accurate results than the optical image-based point cloud (Ganz *et al.* 2019).

From ecosystem to landscape scale: monitoring and management

UAV remote sensing has been used to detect, monitor and fight forest fires (Pastor *et al.* 2011). Equipped with visual, infrared and thermal cameras, UAVs can effectively track fires, predict their expansion and provide real-time information to firefighters

Table 2: Ecological concerns, UAV data requirements and current studies using UAV remote sensing at different ecological scales

Ecological scale	Ecological concerns	UAV data requirement	Previous studies using UAV remote sensing
Individual	The relationship between an individual organism and its environmental factors, including the relationship between individual growth, environmental conditions and biological adaptability to the environment.	<i>Sensors:</i> RGB, multispectral, hyperspectral, LiDAR, thermal <i>Temporal-spatial resolution:</i> day to year; millimeter to decimeter	Individual detection [90, 143, 191, 217, 313, 251, 385, 399]; location [64, 82, 249]; number [69, 99]; growth [201, 250, 400]; crown structure [201, 249]; crown width [64, 249]; crown volume [365]; crown projection area [143]; height [64, 82, 249, 251, 384, 400]; stem diameter [55]; phenology [400]
Population	The variation of population size or quantity in time or space and its regulatory mechanisms.	<i>Sensors:</i> RGB, multispectral, hyperspectral <i>Temporal-spatial resolution:</i> month to year; centimeter to decimeter	Population size [163, 194, 262]; spatial distribution mapping [165, 167, 193, 196, 210, 231, 238, 360]; spatial and temporal variability [130, 164, 181, 275, 351]; seedling emergence [38, 95, 279, 309, 341]; phenological traits [239, 248]
Community	The relationship between community and environment. It mainly studies the structure, function, formation and development of plant community as well as the interrelationship with the environment.	<i>Sensors:</i> RGB, multispectral, hyperspectral, LiDAR, thermal <i>Temporal-spatial resolution:</i> month to year; centimeter to decimeter	Interspecific relationship [3, 37]; functional traits [4, 17, 36, 39, 43, 49, 61, 77, 78, 106, 180, 190, 247, 316]; canopy structure [16, 20, 81, 86, 91, 110, 184, 202, 219, 307, 394]; community structure [31, 65, 70, 75, 87, 94, 105, 111, 126, 282, 377, 379]; phenology [22, 27, 100, 256, 314, 392]; aboveground biomass [23, 57, 63, 131, 135, 138, 148, 152, 170, 182, 188, 218, 220, 258]; communities classification [32, 127, 224, 298, 327]; relationship between canopy variables and biodiversity patterns [40]; community recovery monitoring [42, 237]; community composition [47, 205, 225, 226, 263, 264, 356];

Table 2: Continued

Ecological scale	Ecological concerns	UAV data requirement	Previous studies using UAV remote sensing
Ecosystem	The ecosystem processes, structures, functions, management, material cycles and energy flows.	<i>Sensors:</i> RGB, multispectral, hyperspectral, LiDAR, thermal <i>Temporal–spatial resolution:</i> month to decades; centimeter to decimeter	community disturbances [54, 141, 192]; community health [28, 72, 159, 236, 278, 369, 350]; spatial and temporal variability [74, 204, 207, 223, 355, 358, 368]; biodiversity [102, 333]; effect of canopy structure on light interception [376]
Landscape	The spatial structures, interactions, coordination functions and dynamics of the entirety of many different ecosystems.	<i>Sensors:</i> RGB, multispectral, hyperspectral, LiDAR, thermal <i>Temporal–spatial resolution:</i> month to decades; centimeter to meter	Plant diseases and insect pests [156, 227, 270, 273, 287, 293, 335, 342, 390]; productivity [35, 62, 83, 84, 96, 139, 145, 153, 166, 288, 391, 397]; biological invasion [183, 268, 340, 345]; ecosystem management [185, 265, 269, 276, 318, 397]; ecosystem monitoring [45, 260, 277, 297]; habitat monitoring [68, 80, 132, 304]; disturbance feedbacks [155]; environmental monitoring [53, 59, 119, 178, 254, 322]; fire monitoring [234]; relationship between vegetation structure and the thermal environment [147]; influences of human disturbance [370]
			Habitat fragmentation [48]; land cover classification [108, 154, 169, 187, 1299, 344]; vegetation type classification [109, 136, 198, 200]; farming landscape management [142, 176]; ecosystem phenology [206]

Note: The references can be found in [Appendix S1: Reviewed references list](#).

(Bradley and Taylor 2015; Merino *et al.* 2012). A fire management system on a UAV remote sensing platform was demonstrated in Yuan *et al.* (2015). In addition to fire management, the visual, thermal, multispectral, hyperspectral and LiDAR data acquired from UAV remote sensing can reveal the biotic and abiotic variations of an ecosystem (Valbuena *et al.* 2020). Such data are effective for long-term ecosystem monitoring and management. Mancini *et al.* (2013), for instance, used an SfM image-based approach to generate a DSM of a beach dune system in Italy; the essential features and complexity of the beach dune habitat were shown in the DSM, which provided basic data for future ecosystem management. In addition, the estimation of LAI, nitrogen content, pigment content and water stress via UAV remote sensing can provide a foundation for precise irrigation and fertilization in agriculture (Mathews and Jensen 2013). UAV remote sensing also serves as a cost-effective and flexible tool for monitoring ecological succession (de Almeida *et al.* 2020), restoration (Knoth *et al.* 2013; Zahawi *et al.* 2015) and natural resource management (Inoue *et al.* 2014; Shahbazi *et al.* 2014; Zhang *et al.* 2016).

At landscape scale, UAV remote sensing was mainly used to monitor habitat fragmentation (Yi 2016), land cover change (Cruz *et al.* 2017) and vegetation change (Nguyen *et al.* 2019). Fixed-wing UAV played a more important role in landscape scale studies due to its longer flight time. However, for land cover or vegetation monitoring, imagery with centimeter-level accuracy was helpful but not required and satellite data were sufficient for most landscape studies (Komarek 2020). Considering the distance of remote control (about 5 km), security and difficulty of data processing, UAV is not ideal for ecological studies on landscape scale. Thus, only a limited number of ecological studies using UAV remote sensing were carried out at landscape scale.

CHALLENGES AND PROSPECTS

UAV remote sensing in plant ecology faces a number of challenges.

Challenges in the use of UAV remote sensing systems

Regulatory constraints are major barriers for the use of UAV remote sensing in ecological studies (Allan *et al.* 2015; Rango *et al.* 2009; Werden *et al.* 2015). Stöcker *et al.* (2017) reviewed the current state of UAV regulations worldwide and reported that the

regulations were still preliminary and varied by region due to the rapid emergence of civil UAVs. In some regions, there are no government agencies established in charge of UAV regulation and management, and it is therefore difficult or time consuming to obtain permission to operate a UAV remote sensing system (Vincent *et al.* 2015).

Both the hardware and software of UAV remote sensing systems require improvement. Although more lightweight and smaller sensor systems have become available, they are still expensive. For instance, the Cubert S185 hyperspectral camera (Cubert GmbH, Germany) weighs only 490 g but costs about 88 000 US\$. A UAV LiDAR system generally costs about 120 000–170 000 US\$. It is worth mentioning that prices are dropping. Recently, a new UAV LiDAR system (DJI L1) that integrates a Livox LiDAR module and a mapping camera, has been officially on sale with a price about 12 000 US\$. Although its accuracy has yet to be verified, this low cost system will significantly improve its acceptance and promote the applications of LiDAR in plant ecology. In addition, the integration between UAV platforms and sensors requires improvement. Except for RGB imager, most of the multispectral, hyperspectral and thermal imager are built independent of UAV platform and need an extra GPS module (such as Cubert S185 hyperspectral imager, TC640 thermal imager). Only to UAV with fully integrated sensors, can sensors be triggered through the flight control system. The majority of UAVs, such as DJI M600 PRO do not allow an external device to share GPS information. Therefore, it is still challenging to link the GPS information of UAVs with the collected hyperspectral images, which complicates the data analysis for ecologists (Sha *et al.* 2018).

Challenges in image processing and analysis

Apart from data collection, data processing and analysis represents a main bottleneck of the ecological applications of UAV remote sensing. Compared with conventional aircraft aerial photography, UAV aerial photography is manifested by the low altitude of the flight platform and the small and nonspecialized camera (Whitehead and Hugenholz 2014). The quality of data acquired by UAV depends on types of UAVs and cameras, which is often characterized by small image amplitude, RGB true color and high spatial resolutions. The attitude angle and heading of UAV often produce deviation and result in unstable photo swing angle and overlap due to the influence of air flow and wind direction. In addition, UAVs are

generally equipped with nonmeasurement cameras that requires high-cost processing. The nonlinear optical distortion (such as barrel or pillow distortion) on the image edge brings challenges for image mosaic and analysis (Hardin and Jensen 2011). The processing of UAV data is quite different from that of satellite data, which produces a new demand on data processing software. Due to the small coverage area of a single UAV image, the mosaic workload of orthographic images is significantly higher than that of satellite remote sensing images, which takes up the majority of processing time. For example, the mosaic of 2000 RGB images (each with 8256×5504 pixels) captured by Nikon D850 poses a great challenge for both software and hardware. Even with a high-performance computer, the image mosaic may need 15–20 h.

Image processing software such as Metashape and Pix4d mapper can perform automatic mosaic for high-resolution RGB images. However, it is still challenging to mosaic multispectral or hyperspectral images with low spatial resolutions, small spatial coverages and few image textures if no concurrent GPS data is available. Advanced algorithms such as scale invariant feature transform have been utilized to select matched points between multispectral images and mosaic images. This method was found to be less affected by image scaling and rotation, illumination change and 3D camera view (Lowe 2004; Ren *et al.* 2017). Also, with the increase of spatial and spectral resolutions, image processing becomes quite time consuming. As a result, more efficient algorithms need to be developed.

For data analysis, one challenge is the generality of the models used to estimate plant ecological parameters from UAV remote sensing data. Current studies on remote sensing of plant ecology are data dependent and case specific. The prediction models proposed in these studies are usually not generalizable due to the uncertainties in data collection and processing (especially radiometric correction), and the differences in sampled study areas, acquisition dates or plant species. The physically based method, to some extent, can solve the model transferability issue. This is because the physically based method can simulate the radiative transfer process within plant leaves and canopies under different circumstances (e.g. different leaf biochemical content, canopy structure and viewing geometry). Machine learning approaches have the potential to capture the nonlinear relationship between remote sensing data and vegetation parameters. By combining

machine learning approaches and physically based models, prediction models with both flexibility and transferability could be developed.

There are also many challenges in fusing multisource remote sensing data. One is the coregistration of multisource data, which aims to geometrically align multisource images. In practice, images derived from multisensors tend to have various spatial resolutions (e.g. 30-cm *vs.* 2-m resolutions), georeference accuracies (e.g. 5-cm *vs.* 1-m geometric errors), spectral characteristics (e.g. RGB *vs.* NIR bands), acquisition dates (e.g. early *vs.* late growing seasons) and viewing geometries (e.g. back *vs.* forward scattering angles). All these factors greatly affect the searching for tie points among multisource images. This process in turn determines the coregistration accuracy. When it comes to the UAV-based remote sensing with large amounts of images, an automatic coregistration workflow is often needed. Recently, some progress has been made in this area. For instance, Scheffler *et al.* (2017) developed an open-source Python package 'AROSICS' (Automated and Robust Open-Source Image Co-Registration Software), which enables the automatic coregistration of multisensor satellite images. However, the applicability of this package to UAV-based images that usually have a higher spatial resolution still needs testing.

Another challenge is how to integrate the information derived from multisource remote sensing data. As mentioned above, multisource remote sensing data provide different information on ground objects. For instance, multispectral data can be used to infer the biophysical parameters of plants (e.g. LAI, percent vegetation cover), hyperspectral data to infer the biochemical or physiological parameters of plants (e.g. leaf chlorophyll and nitrogen contents) and LiDAR data to infer the structure parameters of plants (e.g. plant heights, gap fraction). Although most studies have shown the benefits of adding more information into analysis, there is no consensus on the framework of fusing multisource information. Some recent work indicates that machine learning algorithms (e.g. deep convolution neural networks) have the capability of integrating multisource information at different levels (Yao *et al.* 2019).

Automatic species identification

Species classification provides a foundation for assessing many plant community properties, such as community composition, structure and biodiversity. At present, most species classifications via UAV remote sensing require human participation and

interpretation. The accuracy relies on many factors including sensor type, the integration between the UAV platform and the sensor, image resolution, habitat complexity, operator experience and the coordination between ecologists and technologists. With ‘big data analytics’ and machine learning technology, especially coevolutionary neural network algorithms for image processing (Brodrick *et al.* 2019), automatic species identification through UAV remote sensing is becoming increasingly feasible (Jin *et al.* 2018; Sandino *et al.* 2018). If the dataset used for training is ‘big’ enough, computer learning should theoretically generate a satisfactory classification outcome. Crowdsourcing, i.e. the outsourcing of tasks or data collection among a large group of nonprofessionals, has been demonstrated to be an effective approach for big database construction (Minet *et al.* 2017).

The ground-UAV-airplane/satellite multiscale monitoring system

UAV remote sensing bridges the gaps between ground observations and manned aircraft and satellite remote sensing. This bridge makes it possible to answer basic ecological questions across multiple scales. For instance, D’Oleireoltmanns *et al.* (2012) used UAV to acquire the visual images of a soil erosion area in Morocco at 70 and 400 m height and analyzed the distribution, volume and temporal dynamics of gullies at a local scale. Then the authors assessed the mechanism of soil erosion at the sampled

sites and across the entire region by combining UAV remote sensing with satellite images. The multiscale sampling method can also be used to monitor the biodiversity changes at different scales (Gonzalez *et al.* 2020; Guo *et al.* 2016a) and help to detect the form and drivers of biodiversity–ecosystem function relationships across space and time (Williams *et al.* 2021). Under this premise, the research of spatial scaling of ecological stability might also benefit from ground-UAV-satellite monitoring system (Wang *et al.* 2017). Similar applications of a ground-UAV-satellite framework have been reported for precision agriculture (Matese *et al.* 2015; Zecha *et al.* 2013) and land management (Browning *et al.* 2016). Most of the satellite images are available for public research access. A number of platforms, such as Google Earth Engine (GEE), EarthServer, Docker and the Coupled Model Intercomparison Project (CMIP), have been increasingly used in ecological and environmental studies due to the cloud-based geospatial processing capability and the access to a large collection of geospatial datasets such as Landsat and MODIS without requiring downloading and local handling of the images (Baumann *et al.* 2015; Gorelick *et al.* 2017; Liang *et al.* 2020). The higher resolution images from UAVs can serve as ‘ground truth data’ to train and validate the processing of in-cloud images in these platforms, thereby facilitating the construction of a ground-UAV-airplane/satellite multiscale ecological monitoring system.




Figure 4: The connection between UAV remote sensing and ecology. The contents in blue boxes belong to remote sensing science and technology and those in green boxes belong to ecology. The boxes with half blue and half green represent the interdisciplinary parts. UAV remote sensing is suitable for answering ecological questions derived at individual, population, community, ecosystem and landscape scales. In addition to classical ecological questions, there are some novel questions need to be answered, for instance, the relationship between tree crown structure and species competition; the relationship between canopy biochemical feature and environmental change.

From describing ecological phenomenon to answering ecological questions

The majority of the reviewed studies were published on journals in the field of remote sensing with emphasis on remote sensing technology rather than on ecological issues. Only a few papers focused on answering some basic ecological questions, i.e. the relationship between organisms and the relationship between organism and environment (Rissanen *et al.* 2019; Waite *et al.* 2019; Zhang *et al.* 2016; Zhao *et al.* 2020). Most of previous studies have focused on ecological phenomenon descriptions, i.e. retrieving ecological parameters through UAV remote sensing. Furthermore, these studies rarely examined these parameters to answer ecological questions. There is now a strong call to join technological developments with scientific challenges to answer basic scientific questions (Santos *et al.* 2018).

UAV remote sensing opens new possibilities in plant ecology by addressing classical ecological questions at different ecological scales. Ecologists and remote sensing experts should collaborate to determine how data can be collected and analyzed by UAV remote sensing systems to understand ecological processes, such as photosynthesis, nutrient cycling, interspecific relationships and succession (Fig. 4). At the individual scale, crown maps can be linked with environmental variables to explore the adaptive evolution of species and interspecific relationship, such as the relationship between crown shape, environmental factors and interspecies competition. At the community scale, traditional measurements limited to subcanopy/understory can be combined with parameters of overstory captured by UAV to study community composition and structure, including but not limited to canopy structure, plant functional traits and diversity. At the ecosystem scale, maps or parameters derived from UAV remote sensing can be linked with flux tower measurements to investigate ecosystem process and function, especially ecosystem disturbances (Table 2).

Novel methods are needed to fully exploit the use of UAV data

A number of tools or products developed by remote sensors have played a great role in promoting ecological research. These tools or products originate from spaceborne and airborne remote sensing data and generally focus on solving issues on landscape, regional and global scales. With the development of

UAV remote sensing, data and products with ultra-high spatial resolutions have been available for ecological studies at scales from individual to ecosystem. From species identification to community's biophysical and biochemical structure detection, the new research direction has presented new challenges to the development of remote sensing tools or products.

UAVs bring a 'bird's view' of ecosystems to ecologists, and the view is unprecedentedly clear. However, the value of these data is not fully exploited if they are simply used to retrieve the traditional parameters that can be collected on the ground, such as height, DBH or species diversity. UAV remote sensing data include the color, shadow, density and three-dimensional properties of the canopy, but the quantification and analysis of such data with the goal of answering ecological questions remains a challenge. Previous studies have tried to construct canopy chemical assemblies from living vegetation volumes, and geometrical characteristics of canopy gaps in order to better use UAV remote sensing data (Asner *et al.* 2014; Getzin *et al.* 2014). More ecologically meaningful parameters need to be extracted from those data to better understand the canopy surface irregularity and community heterogeneity. In addition to traditional species biodiversity, new metrics need to be developed to include color, biochemical and canopy structure diversity for biodiversity assessment. Furthermore, many interesting ecological phenomena on canopy and community scale such as the crown shape under different species competition intensity can be further understood by ecologists using UAV remote sensing. Data collection and processing of UAV remote sensing is new and complicated for most ecologists, strengthened collaboration between ecologists and remote sensing professionals is needed to promote the application of UAVs in plant ecology and to answer both old and new ecological questions.

CONCLUSIONS

UAV remote sensing bridges the gaps in both scale and resolution between ground observations, conventional manned aircrafts and satellite remote sensing. The maturity of civilian UAV technology is the origin of ecological application of UAVs, and the emergence of SfM photogrammetry promotes the ecological applications of UAV. Mapping, measuring and monitoring of vegetation are three major applications of UAV remote sensing in plant ecology.

From the species to population scale, physiological assessment, species identification and population mapping are the most reported uses of UAV in the literature. Physiological assessment through UAV remote sensing provides a basis for precision ecosystem management. Nevertheless, the ecological applications need to further integrate remote sensing data and ecological process. Species identification and population mapping are the foundation of studies on biodiversity and many ecological processes. The accuracy of species identification in complex habitats could be improved by the integration of big data technology with a machine learning approach. At the community scale, community structure, diversity and biomass are the major concerns of ecologists, but the application of UAVs in these areas are still in early stages. At the ecosystem scale, ecosystem monitoring and management are the primary research fields of interest for UAV application. UAV remote sensing has played an essential role in fighting forest fires, monitoring ecosystem restoration and providing precision crop management. Future applications of UAV remote sensing in plant ecology should deploy the ground-UAV-airplane/satellite multiscale remote sensing system. Most of the ecological applications of UAV remote sensing have been driven by improving the technology rather than answering ecological scientific questions. Close collaboration between ecologists and remote sensing experts is now needed to improve the use UAV remote sensing for resolving ecological questions.

Supplementary Material

Supplementary material is available at *Journal of Plant Ecology* online.

Appendix S1: Reviewed references list and template of reviewed studies.

Appendix S2: A comparative cost analysis of airborne, spaceborne and drone-based remote sensing system and a detailed description of main characteristics of UAV remote sensing case studies in plant ecology.

Figure S1: Number of UAV studies classified by (a) publication year, (b) sensor type, (c) study scale and (d) processing method.

Funding

This research was supported by GDAS' (Guangdong Academy of Sciences) Special Project of Science and Technology Development (2020GDASYL-20200301003, 2017GDASCX-0805, 2020GDASYL-040101, 2020GDASYL-20200102001),

Strategic Priority Research Program of the Chinese Academy of Sciences (XDA13020506), Science and Technology Projects of Guangdong Province (2017A020216022, 2018B030324002) and the National Natural Science Foundation of China (31770473).

Acknowledgements

We thank Professor Bruce Jaffee from University of California at Davis for his advices about technical writing.

Conflict of interest statement. The authors declare that they have no conflict of interest.

Authors' Contributions

Z. Sun and Y. Xie conceived the study; Z. Sun and X. Wang acquired the data; Z. Sun, L. Yang and Y. Huang analyzed the data; Z. Sun wrote the first draft; Z. Wang contributed to the revision of the text. All authors provided input to the final draft.

REFERENCES

- Aasen H, Honkavaara E, Lucieer A, *et al.* (2018) Quantitative remote sensing at ultra-high resolution with UAV spectroscopy: a review of sensor technology, measurement procedures, and data correction workflows. *Remote Sens* **10**:1091.
- Abeyinghe T, Milas AS, Arend K, *et al.* (2019) Mapping invasive *Phragmites australis* in the Old Woman Creek estuary using UAV remote sensing and machine learning classifiers. *Remote Sens* **11**:1380.
- Adam E, Mutanga O, Rugege D (2010) Multispectral and hyperspectral remote sensing for identification and mapping of wetland vegetation: a review. *Wetl Ecol Manag* **18**:281–296.
- Adão T, Hruška J, Pádua L, *et al.* (2017) Hyperspectral imaging: a review on UAV-based sensors, data processing and applications for agriculture and forestry. *Remote Sens* **9**:1110.
- Allan BM, Ierodiaconou D, Nimmo DG, *et al.* (2015) Free as a drone: ecologists can add UAVs to their toolbox. *Front Ecol Environ* **13**:354–355.
- Alvarez-Taboada F, Paredes C, Julián-Pelaz J (2017) Mapping of the invasive species *Hakea sericea* using unmanned aerial vehicle (UAV) and WorldView-2 imagery and an object-oriented approach. *Remote Sens* **9**:913.
- Anderson K, Gaston KJ (2013) Lightweight unmanned aerial vehicles will revolutionize spatial ecology. *Front Ecol Environ* **11**:138–146.
- Ashraf MS, Brabyn L, Hicks BJ, *et al.* (2010) Satellite remote sensing for mapping vegetation in New Zealand freshwater environments: a review. *New Zeal Geogr* **66**:33–43.
- Asner GP, Martin RE, Tupayachi R, *et al.* (2014) Amazonian functional diversity from forest canopy chemical assembly. *Proc Natl Acad Sci U S A* **111**:5604–5609.

- Bagaram MB, Giuliarelli D, Chirici G, *et al.* (2018) UAV remote sensing for biodiversity monitoring: are forest canopy gaps good covariates? *Remote Sens* **10**:1397.
- Baluja J, Diago MP, Balda P, *et al.* (2012) Assessment of vineyard water status variability by thermal and multispectral imagery using an unmanned aerial vehicle (UAV). *Irrig Sci* **30**:511–522.
- Banerjee BP, Raval S, Cullen PJ (2017) High-resolution mapping of upland swamp vegetation using an unmanned aerial vehicle-hyperspectral system. *J Spectr Imaging* **6**:a6.
- Baumann P, Mazzetti P, Ungar J, *et al.* (2015) Big data analytics for earth sciences: the EarthServer approach. *Int J Digit Earth* **9**:1–27.
- Bendig J, Bolten A, Bennertz S, *et al.* (2014) Estimating biomass of barley using crop surface models (CSMs) derived from UAV-based RGB imaging. *Remote Sens* **6**:10395–10412.
- Berni JAJ, Zarco-Tejada PJ, Suarez L, *et al.* (2009) Thermal and narrowband multispectral remote sensing for vegetation monitoring from an unmanned aerial vehicle. *IEEE Trans Geosci Remote Sens* **47**:722–738.
- Berra EF, Gaulton R, Barr S (2019) Assessing spring phenology of a temperate woodland: a multiscale comparison of ground, unmanned aerial vehicle and Landsat satellite observations. *Remote Sens Environ* **223**:229–242.
- Bradley BA (2014) Remote detection of invasive plants: a review of spectral, textural and phenological approaches. *Biol Invasions* **16**:1411–1425.
- Bradley JM, Taylor CN (2015) Georeferenced mosaics for tracking fires using unmanned miniature air vehicles. *J Aersp Comput Inf Commun* **8**:295–309.
- Brodrick PG, Davies AB, Asner GP (2019) Uncovering ecological patterns with convolutional neural networks. *Trends Ecol Evol* **34**:734–745.
- Browning DM, Rango A, Karl JW, *et al.* (2016) Emerging technological and cultural shifts advancing drylands research and management. *Front Ecol Environ* **13**:52–60.
- Calderón R, Navas-Cortés JA, Lucena C, *et al.* (2013) High-resolution airborne hyperspectral and thermal imagery for early detection of Verticillium wilt of olive using fluorescence, temperature and narrow-band spectral indices. *Remote Sens Environ* **139**:231–245.
- Campos-Vargas C, Sanchez-Azofeifa A, Laakso K, *et al.* (2020) Unmanned aerial system and machine learning techniques help to detect dead woody components in a tropical dry forest. *Forests* **11**:827.
- Cao J, Leng W, Liu K, *et al.* (2018) Object-based mangrove species classification using unmanned aerial vehicle hyperspectral images and digital surface models. *Remote Sens* **10**:89.
- Chabot D, Bird DM (2013) Small unmanned aircraft: precise and convenient new tools for surveying wetlands. *J Unmanned Veh Syst* **1**:15–24.
- Chisholmryan A, Cui J, Lumshawn KY, *et al.* (2013) UAV LiDAR for below-canopy forest surveys. *J Unmanned Veh Syst* **1**:61–68.
- Colomina I, Molina P (2014) Unmanned aerial systems for photogrammetry and remote sensing: a review. *ISPRS J Photogramm Remote Sens* **92**:79–97.
- Cruz HO, Eckert M, Meneses JM, *et al.* (2017) Precise real-time detection of nonforested areas with UAVs. *IEEE Trans Geosci Remote Sens* **55**:632–644.
- Cunliffe AM, Brazier RE, Anderson K (2016) Ultra-fine grain landscape-scale quantification of dryland vegetation structure with drone-acquired structure-from-motion photogrammetry. *Remote Sens Environ* **183**:129–143.
- D'Oleireoltmanns S, Marzloff I, Peter KD, *et al.* (2012) Unmanned aerial vehicle (UAV) for monitoring soil erosion in Morocco. *Remote Sens* **4**:3390–3416.
- de Almeida DRA, Zambrano AMA, Broadbent EN, *et al.* (2020) Detecting successional changes in tropical forest structure using GatorEye drone-borne LiDAR. *Biotropica* **52**:1–13.
- Doughty CL, Cavanaugh KC (2019) Mapping coastal wetland biomass from high resolution unmanned aerial vehicle (UAV) imagery. *Remote Sens* **11**:540.
- Dunford R, Michel K, Gagnage M, *et al.* (2009) Potential and constraints of Unmanned Aerial Vehicle technology for the characterization of Mediterranean riparian forest. *Int J Remote Sens* **30**:4915–4935.
- Fletcher AT, Erskine PD (2012) Mapping of a rare plant species (*Boronia deanei*) using hyper-resolution remote sensing and concurrent ground observation. *Ecol Manage* **13**:195–198.
- Floreano D, Wood RJ (2015) Science, technology and the future of small autonomous drones. *Nature* **521**:460–466.
- Flynn KF, Chapra SC (2014) Remote sensing of submerged aquatic vegetation in a shallow non-turbid river using an unmanned aerial vehicle. *Remote Sens* **6**:12815–12836.
- Formoo J, Anderson K, Macleod CJA, *et al.* (2019) Structure from motion photogrammetry in ecology: does the choice of software matter? *Ecol Evol* **9**:12964–12979.
- Ganz S, Kaber Y, Adler P (2019) Measuring tree height with remote sensing—a comparison of photogrammetric and LiDAR data with different field measurements. *Forests* **10**:694.
- Geipel J, Link J, Claupein W (2014) Combined spectral and spatial modeling of corn yield based on aerial images and crop surface models acquired with an unmanned aircraft system. *Remote Sens* **6**:10335.
- Getzin S, Nuske RS, Wiegand K, *et al.* (2014) Using unmanned aerial vehicles (UAV) to quantify spatial gap patterns in forests. *Remote Sens* **6**:6988–7004.
- Getzin S, Wiegand K, Schöning I (2012) Assessing biodiversity in forests using very high-resolution images and unmanned aerial vehicles. *Methods Ecol Evol* **3**:397–404.
- Gil-Docampo ML, Arza-García M, Ortiz-Sanz J, *et al.* (2020) Above-ground biomass estimation of arable crops using UAV-based SfM photogrammetry. *Geocarto Int* **35**:687–699.
- Gomez C, Alejandro P, Hermosilla T, *et al.* (2019) Remote sensing for the Spanish forests in the 21st century: a review of advances, needs, and opportunities. *For Syst* **28**:eR001.
- Gonzalez A, Germain RM, Srivastava DS, *et al.* (2020) Scaling-up biodiversity-ecosystem functioning research. *Ecol Lett* **23**:757–776.
- Gonzalez-Jaramillo V, Fries A, Bendix J (2019) AGB estimation in a tropical mountain forest (TMF) by means of RGB and multispectral images using an unmanned aerial vehicle (UAV). *Remote Sens* **11**:1413.

- Gorelick N, Hancher M, Dixon M, *et al.* (2017) Google Earth Engine: planetary-scale geospatial analysis for everyone. *Remote Sens Environ* **202**:18–27.
- Guerra-Hernandez J, Gonzalez-Ferreiro E, Sarmiento A, *et al.* (2016) Using high resolution UAV imagery to estimate tree variables in *Pinus pinea* plantation in Portugal. *For Syst* **25**:eSC09.
- Guo Q, Liu J, Li Y, *et al.* (2016a) A near-surface remote sensing platform for biodiversity monitoring: perspectives and prospects. *Biodivers Sci* **24**:1249–1266.
- Guo Q, Wu F, Hu T, *et al.* (2016b). Perspectives and prospects of unmanned aerial vehicle in remote sensing monitoring of biodiversity. *Biodivers Sci* **24**:1267–1278.
- Haboudane D, Miller JR, Pattey E, *et al.* (2004) Hyperspectral vegetation indices and novel algorithms for predicting green LAI of crop canopies: modeling and validation in the context of precision agriculture. *Remote Sens Environ* **90**:337–352.
- Haboudane D, Miller JR, Tremblay N, *et al.* (2002) Integrated narrow-band vegetation indices for prediction of crop chlorophyll content for application to precision agriculture. *Remote Sens Environ* **84**:416–426.
- Hardin PJ, Jensen RR (2011) Small-scale unmanned aerial vehicles in environmental remote sensing: challenges and opportunities. *Glsci Remote Sens* **48**:99–111.
- Hassanalian M, Abdelkefi A (2017) Classifications, applications, and design challenges of drones: a review. *Prog Aerosp Sci* **91**:99–131.
- He K, Zhang J, Zhang Q (2009) Linking variability in species composition and MODIS NDVI based on beta diversity measurements. *Acta Oecol* **35**:14–21.
- Hill DJ, Tarasoff C, Whitworth GE, *et al.* (2017) Utility of unmanned aerial vehicles for mapping invasive plant species: a case study on yellow flag iris (*Iris pseudacorus* L.). *Int J Remote Sens* **38**:2083–2105.
- Huang CY, Asner GP (2009) Applications of remote sensing to alien invasive plant studies. *Sensors (Basel)* **9**:4869–4889.
- Hung C, Zhe X, Sukkarieh S (2014) Feature learning based approach for weed classification using high resolution aerial images from a digital camera mounted on a UAV. *Remote Sens* **6**:12037–12054.
- Husson E, Hagner O, Ecke F, *et al.* (2013) Unmanned aircraft systems help to map aquatic vegetation. *Appl Veg Sci* **17**:567–577.
- Husson E, Lindgren F, Ecke F (2014) Assessing biomass and metal contents in riparian vegetation along a pollution gradient using an unmanned aircraft system. *Water Air Soil Pollut* **225**:1–14.
- Huylenbroeck L, Laslier M, Dufour S, *et al.* (2020) Using remote sensing to characterize riparian vegetation: a review of available tools and perspectives for managers. *J Environ Manage* **267**:110652.
- Inoue T, Nagai S, Yamashita S, *et al.* (2014) Unmanned aerial survey of fallen trees in a deciduous broadleaved forest in eastern Japan. *PLoS One* **9**:e109881.
- Ishihama F, Watabe Y, Oguma H (2012) Validation of a high-resolution, remotely operated aerial remote-sensing system for the identification of herbaceous plant species. *Appl Veg Sci* **15**:383–389.
- Jensen J, Mathews A (2016) Assessment of image-based point cloud products to generate a bare earth surface and estimate canopy heights in a woodland ecosystem. *Remote Sens* **8**:50.
- Jin S, Su Y, Gao S, *et al.* (2018) Deep learning: individual maize segmentation from terrestrial LiDAR data using faster R-CNN and regional growth algorithms. *Front Plant Sci* **9**:866.
- Kalacska M, Arroyomora JP, Gea JD, *et al.* (2013) Videographic analysis of *Eriophorum vaginatum* spatial coverage in an ombrotrophic bog. *Remote Sens* **5**:6501–6512.
- Kaneko K, Nohara S (2014) Review of effective vegetation mapping using the UAV (unmanned aerial vehicle) method. *J Geogr Inf Syst* **6**:733–742.
- Karl JW, Yelich JV, Ellison MJ, *et al.* (2020) Estimates of willow (*Salix* spp.) canopy volume using unmanned aerial systems. *Rangeland Ecol Manage* **73**:531–537.
- Keddy PA (2007) *Plants and Vegetation*. Cambridge, UK: Cambridge University Press.
- Klosterman S, Melaas E, Wang J, *et al.* (2018) Fine-scale perspectives on landscape phenology from unmanned aerial vehicle (UAV) photography. *Agric For Meteorol* **248**:397–407.
- Knoth C, Klein B, Prinz T, *et al.* (2013) Unmanned aerial vehicles as innovative remote sensing platforms for high-resolution infrared imagery to support restoration monitoring in cut-over bogs. *Appl Veg Sci* **16**:509–517.
- Komarek J (2020) The perspective of unmanned aerial systems in forest management: do we really need such details? *Appl Veg Sci* **23**:1–4.
- Kwon YS, Pyo J, Kwon YH, *et al.* (2020) Drone-based hyperspectral remote sensing of cyanobacteria using vertical cumulative pigment concentration in a deep reservoir. *Remote Sens Environ* **236**:111517.
- Laliberte A, Craig W, Albert R (2011) UAS remote sensing missions for rangeland applications. *Geocarto Int* **26**:141–156.
- Lary DJ, Alavi AH, Gandomi AH, *et al.* (2016) Machine learning in geosciences and remote sensing. *Geosci Front* **7**:3–10.
- Leduc MB, Knudby AJ (2018) Mapping wild leek through the forest canopy using a UAV. *Remote Sens* **10**:70.
- Leinonen I, Grant OM, Tagliavia CP, *et al.* (2006) Estimating stomatal conductance with thermal imagery. *Plant Cell Environ* **29**:1508–1518.
- Li QS, Wong FKK, Fung T (2017) Assessing the utility of UAV-borne hyperspectral image and photogrammetry derived 3D data for wetland species distribution quick mapping. *ISPRS Int Arch Photogramm Remote Sens Spat Inf Sci* **XLII-2/W6**:209–215.
- Lian PK, Wich SA (2012) Dawn of drone ecology: low-cost autonomous aerial vehicles for conservation. *Trop Conserv Sci* **5**:121–132.
- Liang JY, Xie YC, Sha ZY, *et al.* (2020) Modeling urban growth sustainability in the cloud by augmenting Google Earth Engine (GEE). *Comput Environ Urban* **84**:101542.
- Lopez-Jimenez E, Vasquez-Gomez JI, Sanchez-Acevedo MA, *et al.* (2019) Columnar cactus recognition in aerial images using a deep learning approach. *Ecol Inform* **52**:131–138.
- Lowe DG (2004) Distinctive image features from scale-invariant keypoints. *Int J Comput Vis* **60**:91–110.

- Man Q, Guo H, Shi R (2014) Light detection and ranging and hyperspectral data for estimation of forest biomass: a review. *J Appl Remote Sens* **8**:69–72.
- Mancini F, Dubbini M, Gattelli M, *et al.* (2013) Using unmanned aerial vehicles (UAV) for high-resolution reconstruction of topography: the structure from motion approach on coastal environments. *Remote Sens* **5**:6880–6898.
- Manfreda S, McCabe MF, Miller PE, *et al.* (2018) On the use of unmanned aerial systems for environmental monitoring. *Remote Sens* **10**:641.
- Mariano GC, Morellato LPC, Almeida J, *et al.* (2016) Modeling plant phenology database: blending near-surface remote phenology with on-the-ground observations. *Ecol Eng* **91**:396–408.
- Matese A, Toscano P, Di Gennaro SF, *et al.* (2015) Intercomparison of UAV, aircraft and satellite remote sensing platforms for precision viticulture. *Remote Sensing* **7**:2971–2990.
- Mathews AJ, Jensen JLR (2013) Visualizing and quantifying vineyard canopy LAI using an unmanned aerial vehicle (UAV) collected high density structure from motion point cloud. *Remote Sens* **5**:2164–2183.
- Merino L, Caballero F, Martínez-De-Dios JR, *et al.* (2012) An unmanned aircraft system for automatic forest fire monitoring and measurement. *J Intell Robot Syst* **65**:533–548.
- Messina G, Modica G. (2020) Applications of UAV thermal imagery in precision agriculture: state of the art and future research outlook. *Remote Sens* **12**:1491.
- Minet J, Curnel Y, Gobin A, *et al.* (2017) Crowdsourcing for agricultural applications: a review of uses and opportunities for a farmsourcing approach. *Comput Electron Agric* **142**:126–138.
- Mora C, Vieira G, Pina P, *et al.* (2015) Land cover classification using high-resolution aerial photography in Adventdalen, Svalbard. *Geogr Ann* **97**:473–488.
- Mu Y, Fujii Y, Takata D, *et al.* (2018) Characterization of peach tree crown by using high-resolution images from an unmanned aerial vehicle. *Hortic Res* **5**:74.
- Myneni RB, Ross J (2012) *Photon-Vegetation Interactions: Applications In Optical Remote Sensing And Plant Ecology*. Springer Science & Business Media.
- Nguyen U, Glenn EP, Dang TD, *et al.* (2019) Mapping vegetation types in semi-arid riparian regions using random forest and object-based image approach: a case study of the Colorado River Ecosystem, Grand Canyon, Arizona. *Ecol Inform* **50**:43–50.
- Oliveira RA, Nasi R, Niemelainen O, *et al.* (2020) Machine learning estimators for the quantity and quality of grass swards used for silage production using drone-based imaging spectrometry and photogrammetry. *Remote Sens Environ* **246**:111830.
- Padua L, Vanko J, Hruska J, *et al.* (2017) UAS, sensors, and data processing in agroforestry: a review towards practical applications. *Int J Remote Sens* **38**:2349–2391.
- Pastor E, Barrado C, Royo P, *et al.* (2011) Architecture for a helicopter-based unmanned aerial systems wildfire surveillance system. *Geocarto Int* **26**:113–131.
- Peña JM, Torres-Sánchez J, de Castro AI, *et al.* (2013) Weed mapping in early-season maize fields using object-based analysis of unmanned aerial vehicle (UAV) images. *PLoS One* **8**:e77151.
- Perich G, Hund A, Anderegg J, *et al.* (2020) Assessment of multi-image unmanned aerial vehicle based high-throughput field phenotyping of canopy temperature. *Front Plant Sci* **11**:150.
- Petrich L, Lohrmann G, Neumann M, *et al.* (2020) Detection of *Colchicum autumnale* in drone images, using a machine-learning approach. *Precis Agric* **21**:1291–1303.
- Pichon L, Leroux C, Macombe C, *et al.* (2019) What relevant information can be identified by experts on unmanned aerial vehicles' visible images for precision viticulture? *Precis Agric* **20**:278–294.
- Plesoianu AI, Stupariu MS, Sandric I, *et al.* (2020) Individual tree-crown detection and species classification in very high-resolution remote sensing imagery using a deep learning ensemble model. *Remote Sens* **12**:2426.
- Pölonen I, Saari H, Kaivosoja J (2013) Hyperspectral imaging based biomass and nitrogen content estimations from light-weight UAV. *Proc SPIE Int Soc Opt Eng* **8887**:521–525.
- Rango A, Laliberte A, Herrick JE, *et al.* (2009) Unmanned aerial vehicle-based remote sensing for rangeland assessment, monitoring, and management. *J Appl Remote Sens* **3**:033542.
- Rango A, Laliberte A, Steele C, *et al.* (2006) Using unmanned aerial vehicles for rangelands: current applications and future potentials. *Environ Pract* **8**:159–168.
- Ren JG, Xu P, Yong HL, *et al.* (2017) Ground-to-satellite quantum teleportation. *Nature* **549**:70–73.
- Rissanen K, Martin-Guay MO, Riopel-Bouvier AS, *et al.* (2019) Light interception in experimental forests affected by tree diversity and structural complexity of dominant canopy. *Agr Forest Meteorol* **278**:107655.
- Romero M, Luo Y, Su B, *et al.* (2018) Vineyard water status estimation using multispectral imagery from an UAV platform and machine learning algorithms for irrigation scheduling management. *Comput Electron Agric* **147**:109–117.
- Roosjen PPJ, Suomalainen JM, Bartholomeus HM, *et al.* (2017) Mapping reflectance anisotropy of a potato canopy using aerial images acquired with an unmanned aerial vehicle. *Remote Sensing* **9**:417.
- Rougean JL, Breon FM (1995) Estimating PAR absorbed by vegetation from bidirectional reflectance measurements. *Remote Sens Environ* **51**:375–384.
- Rouse JW, Haas RH, Schell JA, *et al.* (1974) *Monitoring the Vernal Advancement and Retrogradation (Greenwave Effect) of Natural Vegetation*. Greenbelt, MD: NASA/GSFC Type III Final Report, 371.
- Saarinen N, Vastaranta M, Näsi R, *et al.* (2018) Assessing biodiversity in boreal forests with UAV-based photogrammetric point clouds and hyperspectral imaging. *Remote Sens* **10**:338.
- Sanders A (2017) Mapping the distribution of understory *Rhododendron ponticum* using low-tech multispectral UAV derived imagery. In Díaz-Delgado R, Lucas R Hurford C (eds). *The Roles of Remote Sensing in Nature Conservation: A Practical Guide and Case Studies*. Cham, Switzerland: Springer International Publishing, 167–181.

- Sandino J, Gonzalez F, Mengersen K, *et al.* (2018) UAVs and machine learning revolutionising invasive grass and vegetation surveys in remote arid lands. *Sensors* **18**:605.
- Santos MM, Jorge PAS, Coimbra J, *et al.* (2018) The last frontier: coupling technological developments with scientific challenges to improve hazard assessment of deep-sea mining. *Sci Total Environ* **627**:1505–1514.
- Scheffler D, Hollstein A, Diedrich H, *et al.* (2017) AROSICS: an automated and robust open-source image co-registration software for multi-sensor satellite data. *Remote Sens* **7**:676.
- Schneider FD, Kukenbrink D, Schaepman ME, *et al.* (2019) Quantifying 3D structure and occlusion in dense tropical and temperate forests using close-range LiDAR. *Agric For Meteorol* **268**:249–257.
- Schneider FD, Morsdorf F, Schmid B, *et al.* (2017) Mapping functional diversity from remotely sensed morphological and physiological forest traits. *Nat Commun* **8**:1441.
- Sha Z, Wang Y, Bai Y, *et al.* (2018) Comparison of leaf area index inversion for grassland vegetation through remotely sensed spectra by unmanned aerial vehicle and field-based spectroradiometer. *J Plant Ecol* **12**:395–408.
- Shahbazi M, Théau J, Ménard P (2014) Recent applications of unmanned aerial imagery in natural resource management. *GISci Remote Sens* **51**:339–365.
- Singh KK, Frazier AE (2018) A meta-analysis and review of unmanned aircraft system (UAS) imagery for terrestrial applications. *Int J Remote Sens* **39**:5078–5098.
- Smigaj M, Gaulton R, Suarez JC, *et al.* (2017) Use of miniature thermal cameras for detection of physiological stress in conifers. *Remote Sens* **9**:957.
- Solvin TM, Puliti S, Steffenrem A (2020) Use of UAV photogrammetric data in forest genetic trials: measuring tree height, growth, and phenology in Norway spruce (*Picea abies* L. Karst.). *Scand J For Res* **35**:322–333.
- Stöcker C, Bennett R, Nex F, *et al.* (2017) Review of the current state of UAV regulations. *Remote Sens* **9**:459.
- Sun Z, Chen Y, Yang L, *et al.* (2017) Small unmanned aerial vehicles for low-altitude remote sensing and its application progress in ecology. *J Appl Ecol* **28**:528–536.
- Surovy P, Ribeiro NA, Panagiotidis D (2018) Estimation of positions and heights from UAV-sensed imagery in tree plantations in agrosilvopastoral systems. *Int J Remote Sens* **39**:4786–4800.
- Tamouridou AA, Alexandridis TK, Pantazi XE, *et al.* (2017) Evaluation of UAV imagery for mapping *Silybum marianum* weed patches. *Int J Remote Sens* **38**:2246–2259.
- Tian J, Wang L, Li X, *et al.* (2017) Comparison of UAV and WorldView-2 imagery for mapping leaf area index of mangrove forest. *Int J Appl Earth Obs* **61**:22–31.
- Torresan C, Berton A, Carotenuto F, *et al.* (2017) Forestry applications of UAVs in Europe: a review. *Int J Remote Sens* **38**:2427–2447.
- Trichon V (2001) Crown typology and the identification of rain forest trees on large-scale aerial photographs. *Plant Ecol* **153**:301–312.
- Union of Concerned Scientists (2018) <https://www.ucsusa.org/resources/satellite-database>. (15 May 2021, date last accessed).
- Valbuena R, O'Connor B, Zellweger F, *et al.* (2020) Standardizing ecosystem morphological traits from 3D information sources. *Trends Ecol Evol* **35**:656–667.
- Vincent JB, Werden LK, Ditmer MA (2015) Barriers to adding UAVs to the ecologist's toolbox. *Front Ecol Environ* **13**:74–75.
- Waite CE, van der Heijden GMF, Field R, *et al.* (2019) A view from above: unmanned aerial vehicles (UAVs) provide a new tool for assessing liana infestation in tropical forest canopies. *J Appl Ecol* **56**:902–912.
- Wallace L, Lucieer A, Malenovsky Z, *et al.* (2016) Assessment of forest structure using two UAV techniques: a comparison of airborne laser scanning and structure from motion (SfM) point clouds. *Forests* **7**:62.
- Wang Z, Chlus A, Geygan R, *et al.* (2020) Foliar functional traits from imaging spectroscopy across biomes in eastern North America. *New Phytol* **228**:494–511.
- Wang S, Garcia M, Ibrom A, *et al.* (2018) Mapping root-zone soil moisture using a temperature-vegetation triangle approach with an unmanned aerial system: incorporating surface roughness from structure from motion. *Remote Sens* **10**:1978.
- Wang S, Loreau M, Arnoldi JF, *et al.* (2017) An invariability-area relationship sheds new light on the spatial scaling of ecological stability. *Nat Commun* **8**:15211.
- Wang Z, Townsend PA, Schweiger AK, *et al.* (2019a) Mapping foliar functional traits and their uncertainties across three years in a grassland experiment. *Remote Sens Environ* **221**:405–416.
- Wang YY, Zhang K, Tang CL, *et al.* (2019b) Estimation of rice growth parameters based on linear mixed-effect model using multispectral images from fixed-wing unmanned aerial vehicles. *Remote Sens* **11**:1371.
- Watts AC, Ambrosia VG, Hinkley EA (2012) Unmanned aircraft systems in remote sensing and scientific research: classification and considerations of use. *Remote Sens* **4**:1671–1692.
- Werden LK, Vincent JB, Tanner JC, *et al.* (2015) Not quite free yet: clarifying UAV regulatory progress for ecologists. *Front Ecol Environ* **13**:534–535.
- Whitehead K, Hugenholtz CH (2014) Remote sensing of the environment with small unmanned aircraft systems (UASs), part 1: a review of progress and challenges. *J Unmanned Veh Syst* **2**:69–85.
- Whitehead K, Hugenholtz CH, Myshak S, *et al.* (2014) Remote sensing of the environment with small unmanned aircraft systems (UASs), part 2: scientific and commercial applications. *J Unmanned Veh Syst* **2**:86–102.
- Williams LJ, Cavender-Bares J, Townsend PA, *et al.* (2021) Remote spectral detection of biodiversity effects on forest biomass. *Nat Ecol Evol* **5**:46–54.
- Xie Y, Sha Z, Yu M (2008) Remote sensing imagery in vegetation mapping: a review. *J Plant Ecol* **1**:9–23.
- Xie Y, Zhang A, Welsh W (2015) Mapping wetlands and phragmites using publically available remotely sensed images. *Photogramm Eng Rem Sens* **81**:69–78.
- Yang L, Sun Z, Tang G, *et al.* (2016) Identifying canopy species of subtropical forest by lightweight unmanned aerial vehicle remote sensing. *Trop Geogr* **36**:833–839.

- Yang SL, Yang XB, Mo JY (2018) The application of unmanned aircraft systems to plant protection in China. *Precis Agric* **19**:278–292.
- Yao H, Qin R, Chen X (2019) Unmanned aerial vehicle for remote sensing applications—a review. *Remote Sens* **12**:1443.
- Yi SH (2016) FragMAP: a tool for long-term and cooperative monitoring and analysis of small-scale habitat fragmentation using an unmanned aerial vehicle. *Int J Remote Sens* **38**:2686–2697.
- Yin DM, Wang L (2019) Individual mangrove tree measurement using UAV-based LiDAR data: possibilities and challenges. *Remote Sens Environ* **223**:34–49.
- Yuan C, Zhang Y, Liu Z (2015) A survey on technologies for automatic forest fire monitoring, detection, and fighting using unmanned aerial vehicles and remote sensing techniques. *Can J For Res* **45**:783–792.
- Zahawi RA, Dandois JP, Holl KD, *et al.* (2015) Using lightweight unmanned aerial vehicles to monitor tropical forest recovery. *Biol Conserv* **186**:287–295.
- Zarco-Tejada PJ, González-Dugo V, Berni JAJ (2012) Fluorescence, temperature and narrow-band indices acquired from a UAV platform for water stress detection using a micro-hyperspectral imager and a thermal camera. *Remote Sens Environ* **117**:322–337.
- Zecha CW, Link J, Claupein W (2013) Mobile sensor platforms: categorisation and research applications in precision farming. *J Sens Sens Syst* **2**:51–72.
- Zhang Z, Flores P, Ighathinathane C, *et al.* (2020) Wheat lodging detection from UAS imagery using machine learning algorithms. *Remote Sens* **12**:1838.
- Zhang J, Hu J, Lian J, *et al.* (2016) Seeing the forest from drones: testing the potential of lightweight drones as a tool for long-term forest monitoring. *Biol Conserv* **198**:60–69.
- Zhang C, Kovacs JM (2012) The application of small unmanned aerial systems for precision agriculture: a review. *Precis Agric* **13**:693–712.
- Zhao LC, Wang JW, Zhao CZ, *et al.* (2020) Using unmanned aerial vehicles to quantify spatial patterns of *Nitraria tangutorum* and *Reaumuria songarica* shrubs under different sand burial conditions in the Jiayuguan national wetland, northwest China. *Int J Remote Sens* **41**:19–30.
- Zou XD, Liang AJ, Wu BZ, *et al.* (2019) UAV-based high-throughput approach for fast growing *Cunninghamia lanceolata* (Lamb.) cultivar screening by machine learning. *Forests* **10**:815.
- Zweig CL, Burgess MA, Percival HF, *et al.* (2015) Use of unmanned aircraft systems to delineate fine-scale wetland vegetation communities. *Wetlands* **35**:303–309.