



Ailanthus altissima. © Jean-Marc Dufour-Dror.

The management of tree of heaven (*Ailanthus altissima*)

Measures and associated costs

Species (scientific name)	<i>Ailanthus altissima</i> (Mill.) Swingle
Species (common name)	Tree of heaven
Author(s)	Giuseppe Brundu (Department of Agriculture, University of Sassari, Italy)
Date Completed	06/11/2017
Reviewer	Ingo Kowarik (Department of Ecology, Technische Universität Berlin, Germany)
Reference	Brundu, G. 2017. Information on measures and related costs in relation to species considered for inclusion on the Union list: <i>Ailanthus altissima</i> . Technical note prepared by IUCN for the European Commission.

Table of contents

	Summary of the measures	2
	Prevention	3
	Ban on importing	3
	Public awareness raising campaigns	5
	Awareness campaigns	7
	Early detection	9
	Surveying	9
	Rapid eradication	12
	Hand pulling	12
	Management	14
	Integrated control strategy	14
	Classical biological control	17
	Bibliography	19
	Appendix	22

Common names

BG	Айлант
HR	Žljezdasti pajasen
CZ	Pajasan žláznatý
DA	Skyrækker
NL	Hemelboom
EN	Tree of heaven
ET	Näärmeline jumalapuu
FI	Haisujumaltenpuu
FR	Ailante glanduleux
DE	Götterbaum
EL	Αείλανθος ο υψηλότατος
HU	Mirigyés bálványfa
IE	Crann na nDéithe
IT	Ailanto (albero del paradiso)
LV	Augstākais ailants
LT	Aukštasis ailantas
MT	Ix-xumakk il-falz
PL	Bożodrzew gruczołowaty
PT	Espanta-lobos
RO	Cenușer
SK	Pajaseň žliazkatý
SL	Veliki pajesen
ES	Ailanto (árbol del cielo)
SV	Gudaträd



Summary of the measures, emphasizing the most cost-effective options.

Ailanthus altissima (tree of heaven), Simaroubaceae, is an early successional tree, native to China and North Vietnam, introduced as ornamental and for other uses which has become invasive in Europe and on all other continents except Antarctica. It is most abundant in urban habitats and along transportation corridors, but can also invade natural habitats. The measure that can prevent intentional introductions of the species is a ban on keeping, importing, selling, breeding and growing. To evaluate the acceptability of such a ban on keeping, importing, selling, breeding and growing *Ailanthus*, it is important to take into account the actual purposes for *Ailanthus* voluntary introduction and use in the European Union.

To prevent unintentional introduction and spread, the following measures are recommended: (1) identification of large trees (female or male) for eradication; (2) applying best construction and management practice for roads and railways; (3) applying best management practices (for example, limiting disturbance) in habitats and land uses that are at risk of invasion; and (4) awareness campaigns to prevent mislabeling, dumping of garden waste, soil movements from infested sites. As for most of the invasive alien species that have many unintentional pathways of accidental introduction and secondary spread, all the possible measures should be included in a single action plan.

Early detection could be achieved by surveying the highest priority roads and railroads, urban and peri-urban areas, historical and archaeological buildings, riparian network, and natural and semi natural areas crossed or in close proximity. Surveys can be done scouting on foot, by car, by

aerial vehicle (helicopter) and assisted with distal or proximal remote sensing tools such as unmanned aerial vehicles or systems. The measures to achieve rapid eradication of *Ailanthus* are the same as for the species Management, for example, rapid eradication should follow an integrated control methodology.

The management of *Ailanthus* needs to make use of an integrated control strategy within a dedicated management plan. In the case of *Ailanthus* integrated control, specific tactics can include any combination of the following: herbicide rotation and herbicide mixtures, herbicide application rate and timing, precision application (for example, "drill-fill" or stem injection), selection and integration of different methods according to the location and site accessibility, age, size, and the sex of the targeted *Ailanthus* plant (hand pulling can be an option, even if in very few cases), seed bank management, habitat manipulation (both for prevention and control) which can include several management and silvicultural options and disturbance mitigation. Importantly, as studied in the US, rodents can have a significant effect on the survival of tree seeds and seedlings invading old fields.

A variety of insects and diseases affect tree-of-heaven with varying success; however, there are no biological control agents currently approved for use in the United States. In addition, the release of macro-organisms as biological control agents is currently not regulated at EU level. Nevertheless national/regional laws are to be respected. Before any release of an alien species as a biological control agent an appropriate risk assessment should be made.

Measures for preventing the species being introduced, intentionally and unintentionally.

This section assumes that the species is not currently present in a Member State, or part of a Member State's territory.



A ban on importing (pre-border measure), selling, breeding, growing, and cultivation, as required under Article 7 of the IAS Regulation, targeting intentional introduction of plants and propagules of *A. altissima*.

EFFECTIVENESS OF MEASURE

When addressing an alien plant species that is introduced mostly intentionally (such as ornamentals) and not yet present in the territory of the European Union, a ban on keeping, importing, selling, breeding, and growing the species is expected to be an effective measure against invasion. However, *Ailanthus* is already present in most of the Countries of the European Union, therefore such a measure is likely to only limit further spread and re-invasion in sites where removal or control intervention are taking place. In addition, *Ailanthus* spread also occurs unintentionally (cf. the following *Prevention* sections).

EFFORT REQUIRED

To be effective, these restrictions and trade bans must be enforced indefinitely.

RESOURCES REQUIRED

Evaluations of the costs associated with the implementation of a ban on keeping, importing, selling, breeding and growing for *Ailanthus* are not available. However, if the ban for *Ailanthus* will be part of general biosecurity policy and biosecurity strategy, resources and costs will be reduced. For example, if there will be a unique biosecurity strategy for all the invasive alien plants of Union Concern this will produce general beneficial effects and economies of scale, including for the training of staff and application of custom controls as some pathways are responsible for the introduction of more than one taxon.

However, the costs of compliance with such measures are often asymmetrical because compliance depends on technical know-how, production facilities, vectors safety and an infrastructural base that, while usually available in developed and emerging markets, is often lacking in many low-income countries (Murina and Nicita, 2017). As a result, the cost of compliance and the resource required might be different across EU Member States, for example in relation to the existing organisational framework, the total number of points of entry (unpacking facilities, car import yards and

industrial premises where machinery is unloaded), the size of the borders, the size of the country, the total number of islands, the biogeographical region, and the trade routes.

SIDE EFFECTS

The pollen of *Ailanthus* is considered as a possible allergenic source (for example, Ballero *et al.*, 2003) so that a ban on keeping, importing, selling, breeding and growing *Ailanthus* is likely to have a positive effect on human health.

ACCEPTABILITY TO STAKEHOLDERS

To evaluate the acceptability of a ban on keeping, importing, selling, breeding and growing *Ailanthus*, it is important to take into account the many purposes for *Ailanthus* intentional introduction and use in the European Union.

Ailanthus was originally introduced in Europe (18th century) as an amenity tree and as forage for silk worms. Its use as an ornamental and shade tree however continues today in Europe. Further uses include plantations for erosion control on slopes, verges of traffic lanes, dunes on the coast, afforestation or reforestation, for the reclamation of landfill sites and mine spoils (Kowarik and Säumel, 2007, and references cited therein). It grows on a wide range of soils, from poor to rich and from dry to moist. *Ailanthus* tolerates relatively high levels of air pollution and may be able to sequester some pollutants. For this reason, it has been widely planted in urban areas worldwide to reduce environmental pollution (Baptista *et al.*, 2014). *Ailanthus* has been used also for biomass production for fuel wood and for the production of fodder for goats and cattle (Feret, 1985; Baptista *et al.*, 2014).

In order to meet future demand and to overcome the wood shortages, studies have been conducted worldwide to evaluate the potential of new or alternative resources as raw material components for pulp and paper production. Thus, the use of fast growing species such as *Ailanthus* is considered as a promising alternative. The results achieved by Baptista *et al.*, (2014) showed that the properties of the



Ailanthus altissima. © Wolfgang Rabitsch

paper obtained from *Ailanthus* are close to those of the reference ones (*Eucalyptus globulus*).

Moreover, *Ailanthus* has been studied from the point of view of its biological activities and pharmacological applications, proving its use in traditional Chinese medicine and its potential application in modern medicine (Baptista *et al.*, 2014, and references cited therein; Peng *et al.*, 2017). However, *Ailanthus* is traditionally used also in the invaded range, such as for flower tips for vermifuge and antiseptic properties in Sicily, Italy (Napoli, 2008).

The extract or semi-purified fractions of *Ailanthus* are strong plant growth inhibitors, therefore good candidates as potential environmentally safe and effective agricultural pest management agents (Tsao *et al.*, 2002). In addition, *Ailanthus* functions as food for honeybees; the honey is tasty but initially bad smelling (Kowarik and Säumel, 2007, and references cited therein).

However, most of the uses and benefits provided by *Ailanthus* described above can be provided by alternative native woody species, such as in the case of slope stabilisation and reduction of soil erosion risk, shade and other landscaping uses. The economic impact on the ornamental plant industry is likely to be low, as only small volumes of the species are traded. In addition, in the invaded range, the negative effects of *Ailanthus* on ecosystem services are generally considered far greater than positive effects (for example, Andreu *et al.*, 2009; Conedera *et al.*, 2014). It is important

to take into account the potential role of *Ailanthus* in urban systems in providing cultural ecosystem services. As it often grows well in harsh urban-industrial environments where native species may be limited, regulating ecosystem services might be also relevant (see, for example, Kowarik and Säumel, 2007). However, conflicting opinions do exist and would need further clarifications, so that further research and an adequate informative action may convince stakeholders.

Importantly, in some European countries or regions, the legislation in force poses limitations to the use of *Ailanthus*. For example, In Italy, *Ailanthus* is included in a black list for two regions, Lombardy and Piedmont (respectively on the basis of the Lombardy regional law 31 March 2008, No. 10; Piedmont DGR n. 23-2975/2016). In addition, the presence of *Ailanthus* outside cultivation, is generally perceived as negative by many stakeholders (as explained in the sections on unintentional introduction and spread).

ADDITIONAL COST INFORMATION

The exact costs are unknown. However, it is widely accepted that prevention (a ban on keeping, importing, selling, breeding and growing *Ailanthus*) would be highly cost-effective, being much cheaper than early detection, rapid eradication or management measures (cost of inaction).

General information (not specifically concerning *Ailanthus*) can be achieved through the documents and reports of those Countries that have national biosecurity policies in force, such as Australia¹ and New Zealand. Another source of general information on prevention cost is the study of Epanchin-Niell (2017).

LEVEL OF CONFIDENCE*

High.

Ailanthus is unlikely to be confused with other species. When a ban on keeping, importing, selling, breeding and growing an alien species not yet present in the territory of the European Union is addressing an alien plant species that would be introduced mostly intentional such as *Ailanthus*, it can be expected to be an effective measure against invasion. In conclusion, a ban on keeping, importing, selling, breeding and growing will be very effective (with a high level of confidence) to limit the intentional introductions of *Ailanthus* to areas of the EU where the species is presently absent (such as Denmark), where it has been locally eradicated or where eradication attempts are occurring (for example, in the framework of LIFE projects).

1 [http://www.parliament.wa.gov.au/publications/tabledpapers.nsf/displaypaper/4010213a4c3d7d22ffde7bb64825812300165804/\\$file/213.pdf](http://www.parliament.wa.gov.au/publications/tabledpapers.nsf/displaypaper/4010213a4c3d7d22ffde7bb64825812300165804/$file/213.pdf)
<http://www.gja.org.nz/Portals/79/Content/Documents/Resource-Library/GIA%20Annual%20Report%202016-web.pdf?ver=2016-11-21-145322-423>

* See Appendix



General considerations.

MEASURE DESCRIPTION

The measures that can prevent unintentional introduction and spread, should be based on a comprehensive analysis of the pathways of unintentional introduction and spread of *Ailanthus* within the territory of the European Union, and identify the pathways which require priority action.

This section reviews the available knowledge on measures that could prevent or reduce the unintentional introduction and spread of *Ailanthus*. These preventive measures include the application of best management practices for the construction and management of roads and railways, and for habitats and land uses that are at risk of invasion (discussed in this section). In addition, awareness campaigns to prevent mislabeling, dumping of garden waste, soil and seed movements from infested sites and targeting key stakeholder groups will be needed (discussed in the following *Prevention* section). When it comes to control and management measures it is generally agreed (for example, USDA, 2014) that large female individuals should be considered a priority for control, aiming to reduce the risk of further unintentional spread of *Ailanthus* from infested areas to uninvaded areas through seed rain. It is important to note that all these measures addressing unintentional introductions and secondary spread, need to be addressed within a **single action plan**.

Unintentional introductions – Applying best management practice for construction and maintenance of roads and railways, and for habitats and land uses that are at a high risk of invasion.

Ailanthus can disperse at the local scale by seeds and vegetatively. Long-distance dispersal of seeds and clonal ramets is mediated by natural corridors (rivers, see Kowarik and Säumel, 2008; Cabra-Rivas *et al.*, 2014), and by road/rail transport and other human activities and infrastructures. In southern France, Kowarik (1983) observed clonal roadside populations of *Ailanthus* up to a length of 120 m. Clonal growth starting on roadsides encroached into evergreen shrub communities out to a distance of 25 m, and arable fields up to 45 m (Kowarik and Säumel, 2007, and references cited therein). Wind moves samaras both individually as well as aggregated in clusters. Wind was shown to move samaras at least 200 m over a hayfield, with four 18-m-tall trees as seed sources (Landenberger *et al.*, 2007). In urban habitats, sealed surfaces of pavements or streets may facilitate secondary dispersal of samaras by wind along linear habitats (Kowarik and Säumel, 2007 and references cited therein). The seed shadow from an 8-m tall tree extended for 456 m along the sidewalk in an

urban transportation corridor, as a combined function of primary and secondary wind dispersal (Kowarik and von der Lippe, 2011).

Outside of cities, *Ailanthus* preferentially colonizes transportation corridors, and among these mostly road and railroad verges and medians of motorways. Samaras can be attached to vehicles, and are also dispersed by the airflow of cars (von der Lippe *et al.*, 2013). Mainly starting from roadside verges, *Ailanthus* can invade borders of agricultural fields, meadows, vineyards, olive groves and old fields (Kowarik and Säumel, 2007 and references cited therein). Merriam (2003) suggested that the reason that *Ailanthus* is most closely associated with railroad rights-of-way in North Carolina (USA) is because winds from passing trains carry seeds along the tracks more effectively than in other edge habitats.

Hydrochory is a secondary dispersal vector in *Ailanthus* as supported by a study by Kowarik and Säumel (2008). They demonstrated that samaras were able to drift for at least 20 days on water. Both floating and submerged seeds were able to germinate after differing periods of exposure to water. In addition, a significant portion of buried stem fragments developed both adventitious shoots and roots after a stay of 3 or 10 days in water. In a river, water moved a quarter of exposed samaras over 1,200 m in 3 hours (Säumel and Kowarik, 2010).

Agricultural machinery accidental transport of *Ailanthus* seeds has also been documented (for example, Fryer, 2010). Rodents may occasionally act as dispersal agents by padding out their dens with collected samaras. In Texas, several bird species feed on *Ailanthus* seeds (Kowarik and Säumel, 2007, and references cited therein).

However, in the family *Simaroubaceae* we can expect the presence of andromonoecious (male and hermaphrodite flowers occur separately on the same plant), functionally dioecious, or distinctly dioecious plants (male and female reproductive structures on separate plants.). *Ailanthus* is a dioecious tree. Female flowers may have stamina, but these do not contain pollen. Other authors suggested that flowers might also be bisexual or the trees monoecious (male and female reproductive structure are found on the same plant but in different flowers), but empirical data on the occurrence and proportion of flower types in populations are scarce. Possibly, some authors address female flowers with sterile stamina as bisexual or 'hermaphrodite' (Kowarik and Säumel, 2007, and references cited therein). Importantly, in the city of Prague (Czech Republic), 614

mature individuals were examined: 50% of them were male plants with staminate flowers only, 48.5% were female plants bearing pistillate flowers and 1.5% of the trees were bisexual individuals classified as andromonoecious (poster communication at the 2014 NeoBiota Conference, Holec *et al.*, 2014).

There are many guidelines available for **best management practices** in road and railway building and maintenance that help prevent the spread of invasive plants on roadsides and into agricultural or natural areas (for example, Pennsylvania Department of Transportation, 2014; Graziano and Clayton, 2017; USDA Forest Service San Dimas Technology, 2017; For railway see: ÖBB-Infrastruktur AG Guidelines, Austria²). Importantly, activities such as mowing, grading, ditching and construction can work to either exacerbate or prevent the spread of invasive plants, including *Ailanthus*

Those habitats and land uses that are more prone to be invaded by *Ailanthus* should be managed according to **specific guidelines** that should indicate adequate forest management measures, maintenance interventions for transport corridors and urban-forest interfaces, management of riparian networks (Cabra-Rivas *et al.*, 2014), measures to reduce or to contrast land abandon in agricultural and forest areas. This is particularly important as *Ailanthus* is already present and established in many MSs, so that there is a higher risk of accidental introduction and spread into these habitats from infested areas.

Importantly, forest disturbance may provide greater opportunity for invasion of *Ailanthus*, thereby altering the successional trajectory of native plant communities. Forest disturbance is typically characterized by biomass removal that creates new growing space, such as fires, removal of litter, clear cuttings, coppicing, and opening of roads. Rapid establishment and growth, and vegetative reproduction in high light environments make disturbed areas such as timber harvests particularly prone to invasion by *Ailanthus* (Kota *et al.*, 2007; Radtke *et al.*, 2013; Wunder *et al.*, 2016; Wickert *et al.*, 2017). In forest and natural areas, one additional important prevention measure is to wash vehicles before they enter a “weed-free” area or when they leave an *Ailanthus* infested area. A prototype of portable vehicle washer is described in the web site of USDA Forest Invasive Species Program (<https://www.fs.fed.us/invasivespecies/prevention/>). This measure is of course advisable to reduce the accidental transport of many other weeds.

In addition, *Ailanthus* colonizes a broad array of urban habitats ranging from walls, fence rows, cracks of sidewalks, and road and railroad embankments to abandoned lots and

urban parks (Kowarik and Säumel, 2007, and references cited therein), archaeological sites (Celesti-Grapow and Blasi, 2004). The abandonment of these habitats/land, or an inadequate intensity of maintenance interventions, will very likely promote the colonisation by *Ailanthus* and the establishment of large clones.

In Southern Europe, *Ailanthus* can frequently colonise abandoned agricultural fields (Marco *et al.*, 2010), such as abandoned olive groves and vineyards, particularly on limestone areas. The colonisation of abandoned chestnut orchards by *Ailanthus* in Switzerland is reported by Knüsel *et al.*, (2015).

In general, any intervention in archaeological site management³ (including exclusion of domestic herbivores) can enhance the risk of invasions by weeds including *Ailanthus*.

EFFECTIVENESS OF MEASURE

To be effective, the different preventive measures must be integrated into a single action plan and enforced addressing all possible unintentional pathways.

EFFORT REQUIRED

The measures need to be maintained indefinitely.

RESOURCES REQUIRED

No information available. However, general information on the cost of awareness raising campaigns for invasive alien plants can be found in LIFE projects on invasive alien plants.

SIDE EFFECTS

Ailanthus can have negative effects on human infrastructures, including those building of historical and cultural importance, so that preventing its spread is likely to have a positive effect on local and national economies. The pollen of *Ailanthus* is considered as a possible allergenic source (for example, Ballero *et al.*, 2003) so that preventing its spread is likely to have a positive effect on human health.

Applying best construction and management practices for roads and railways will also limit the spread of other invasive alien species with similar ecological requirements and pathways of spread. In addition it has been remarked for the south of Italy that road-side removal of *Ailanthus* might improve road safety due to enhanced visibility of road signals (Casella and Vurro, 2012).

ACCEPTABILITY TO STAKEHOLDERS

Please see the sections on preventive measures for intentional introduction. Applying best construction and

2 http://events.uic.org/IMG/pdf/session_7b_-thomas_schuh_-_obb_-_invasive_alien_plants_.pdf
http://botany.uibk.ac.at/neophyten/download/05_OeBB_Schuh_Broschuere.pdf

3 <http://excellence.minedu.gov.gr/thales/en/thalesprojects/380237>
<http://www.doc.govt.nz/documents/science-and-technical/sap243entire.pdf>

management practices for roads and railways, may incur costs for those sectors required to undertake the measures. However, measures to prevent accidental introduction and secondary spread are more likely to be accepted by stakeholders and to be less conflictual, as *Ailanthus* invasions are generally perceived as a problem in the habitats and land uses where *Ailanthus* stands are established and costly control intervention are often occurring. Importantly, in some European countries or regions, the legislation in force poses limitations to the use of *Ailanthus* and some funding for the control of *Ailanthus* in agricultural areas are available through the EU's rural development policy 2014–2020.

ADDITIONAL COST INFORMATION

No information available.

LEVEL OF CONFIDENCE*

High.

There are a number of best practice guidance documents published. Prevention is the first and most cost-effective line of defence against invasive alien species. In conclusion, the preventive measures summarised in the present section and described in the following sections (with a high level of confidence due to the fact that *Ailanthus* is quite a well-studied species) are very likely to limit the spread of *Ailanthus* to areas of the EU where the species is presently absent or where it has been locally eradicated. Road and railways are very well-known transport corridors for *Ailanthus* accidental spread.



Ailanthus altissima. © Jean-Marc Dufour-Dror



Awareness campaigns.

MEASURE DESCRIPTION

This set of preventive measures can be adopted by EU MSs making use of specific national legislation tools or can be included in more general biosecurity policy and biosecurity strategy for larger groups of invasive alien species. However, in addition to regulations MSs or single stakeholders' categories may consider and use a voluntary code of conduct as an effective alternative or complementary approach (for example, EPPO Phytosanitary Procedures, PP 3/74 (1); EPPO, 2009). Additional information can be found in the EPPO Guidelines for the management of plant health risks of bio-waste of plant origin (EPPO, 2008). Concerning the cleaning of machinery or of other vectors, useful information can be found from the guidelines prepared by Biosecurity Queensland, part of the Department of

Agriculture, Fisheries and Forestry, in Australia (Biosecurity Queensland, 2014) and similar documents⁴.

Simple measures, such as cleaning of agricultural machinery cannot be considered as “stand alone” measures but must be integrated in more general action plans and codes of conduct, and widely disseminated and integrated with other general biosecurity measures (such as the quality of forest and horticultural reproductive materials and nursery plants weed-free).

EFFECTIVENESS OF MEASURE

To be effective, preventive measures must be enforced addressing all possible unintentional pathways.

* See Appendix

⁴ See also: <http://dpiw.tas.gov.au/Documents/Washdown-Guidelines-Edition-1.pdf>
<https://www.boprc.govt.nz/media/395661/keepitclean.pdf>

EFFORT REQUIRED

The measures need to be maintained indefinitely, to prevent further spread of this alien plant within the EU.

RESOURCES REQUIRED

No information available. However, there are many LIFE projects that can provide information on awareness campaign concerning other invasive alien plants. Examples of LIFE projects specifically dealing with *Ailanthus* are reported in the *Management* section below.

SIDE EFFECTS

No information available.

ACCEPTABILITY TO STAKEHOLDERS

See the sections on the preventive measures for unintentional introduction and spread.

The overall objective of the LIFE AlterIAS project was to reduce the introduction of invasive plants at source, by raising awareness about their environmental risks amongst the whole ornamental horticulture supply chain in Belgium. The project aimed to promote best practices for preventing the release and spread of invasive alien species through a voluntary Code of Conduct produced with the involvement of the horticultural sector (Halford *et al.*, 2014).

ADDITIONAL COST INFORMATION

No information available.

LEVEL OF CONFIDENCE*

High.

A number of public awareness campaigns have been ran and codes of conducts published. Prevention is the first and most cost-effective line of defence against invasive alien species.

* See Appendix

Measures for early detection of the species and to run an effective surveillance system to detect efficiently new occurrences.



Surveying.

MEASURE DESCRIPTION

The measures to achieve early detection and run an effective surveillance system for achieving an early detection of a new occurrence of *Ailanthus* should take into account the pathways of introduction and spread, the location and distribution of existing infested areas, the susceptibility to be invaded by the diverse habitats and land uses. Early detection and rapid eradication (EDRE) are critical for preventing establishment of *Ailanthus*. Coordination efforts should be made between land managers, the local public (citizen-science), and road crews, etc., on identification of *Ailanthus* so suspected infestations can be reported.

There is not a single method that can be used, so it is advisable to frame the available measures and options in a dedicated **action plan**.

Early detection can be achieved by **surveying the highest priority roads and railroads, urban and peri urban areas, historical and archaeological buildings, riparian network, and natural and semi natural areas crossed or in close proximity** (at least within 2 km of known locations of *Ailanthus* according to USDA, 2014). Surveys can be done on foot, by car (see McAvoy *et al.*, 2012) or aerial vehicle (helicopter) and assisted with distal or proximal remote sensing tools such as unmanned aerial vehicles or systems (UAV, UAS, drones).

An example of road survey ('windshield survey') methodology is found in McAvoy *et al.*, (2012) for Virginia (USA). The authors state that "young tree-of-heaven stems quickly exceed heights over 1 m and are easily identified by their long compound leaves. The number of living tree-of-heaven stems up to 25 m from the edge of both sides of the road and in the median, if present, were counted and recorded in 1.6-km road segments. At the end of each 1.6-km road segment the total number of tree-of-heaven was recorded and a new count was begun for the next 1.6-km road segment. To determine the amount of error in the windshield counts 10 1.6-km road segments infested with tree-of-heaven were surveyed from the vehicle. At the end of each 1.6-km segment, both sides of the road were walked and the number of tree-of-heaven stems was counted."

Although not specifically planned for *Ailanthus*, Harris *et al.*, (2001) provide guidance and a model for New Zealand on time intervals for active weed and invasive alien plants surveillance and they distinguish active surveillance from fortuitous surveillance. Several factors determine site invasiveness and probability of detection: rate of spread of the invasive alien plant, ability to find the new weed. The rate of arrival of an alien plant at a site varies with the proximity of the site to roads, towns and adjoining land use. Once an alien plant has arrived, its rate of spread depends on the habitat, the growth form, and its inherent biological capacity for spread. The visibility of an alien plant or small population, and hence the probability of finding it, varies with its growth stage, growth form and the location of the infestation. The ability to find an alien is also a function of our search effort (Harris *et al.*, 2001).

In the Czech Republic, a research project (www.invaznirostliny.cz/en) aimed at developing an innovative method of mapping *Ailanthus* and other invasive plant species featuring a dedicated unmanned aerial system (UAS). This project aims to develop an unmanned aircraft system and methodological workflow enabling fast and precise monitoring of invasions. Different approaches to image pre-processing are tested, choosing the best methods for the data acquisition and their geometric and radiometric correction.

In the USA, Rebbeck *et al.*, (2015) tested a digital aerial sketch-mapping system (DASM - <https://www.fs.fed.us/eng/rsac/invasivespecies/documents/DASM.pdf>) run on a laptop computer with a touch screen display and stylus, allowing the manual recording of sketched features onto a base map. The goal of this method was to capitalize on conspicuous seed clusters of *Ailanthus* and test whether application of DASM technology from helicopters could be used as a fast and cost-effective tool to map seed-bearing *Ailanthus* in forested landscapes. These tests were conducted across 289,000 acres of public and private forestlands. Field crews assessed the accuracy of the aerial surveys through ground reference data collection. Crews also conducted aerial surveys during the leaf-on season to determine whether *Ailanthus* foliage and seed clusters were distinguishable

from those of other species. Finally, a mapping protocol was developed for use by the forest management community. The georeferenced data could be used as a management strategy to identify high priority areas to target the removal of *Ailanthus* seed sources to minimize its dispersal and slow its spread (cf. also, http://www.edc.uri.edu/atmt-dss/report_forecast/toh_habitat.html).

Citizen science can also play a role in the early detection, for example the Invaders of Texas Citizen Scientist Program (USA) is designed to train volunteer citizen scientists to detect invasive plants in their area and report them to a state-wide detection and mapping database housed on the www.Texasinvasives.org website (including *Ailanthus*). Another example, is the SISSI citizen science project in Italy (<http://sissi.divulgando.eu/specie/dettaglio/3>) and the LIFE project CSMON (http://www.csmon-life.eu/pagina/dettaglio_specie/35).

EFFECTIVENESS OF THE SURVEILLANCE

Ailanthus can be easily identified in all the stages of its life cycle, so that surveillance is expected to be effective. However, surveillance measures must address all possible areas where *Ailanthus* entry, establishment and spread may occur, giving priority to habitats and land uses more prone to invasion.

Data collected through citizen science need to be carefully screened to avoid false-positives. In addition, surveillance will be facilitated whenever access to private properties will be necessary.

There is not a single method or tool that can be used “stand alone”, so it is advisable to frame the available measures and options in a dedicated action plan. For example, UAV operation in urban and inhabited areas might be prohibited or limited in some areas/MSs, limiting the application especially for invasive species like *Ailanthus* that spread into man-made habitats. Despite its limits, drones still provide a reasonable alternative to satellite imagery especially in vegetation mapping, where often data of high spatial and temporal resolution are required. In addition, they might be of great help in areas that cannot be surveyed on foot.

EFFORT REQUIRED

The measures need to be maintained indefinitely. Early detection of *Ailanthus* in the EU needs availability of well trained staff, to conduct surveillance by vehicle, by helicopter, on foot or by boat. The surveillance done by the trained staff could be supported by non-governmental organisations and “citizen science” activities. In addition, remote sensing techniques, including drones and GIS modelling, may greatly help surveillance.

The road or ‘windshield’ survey undertaken by McAvoy *et al.*, (2012) took place along the secondary, primary,

and interstate roads from May to September in 2004, 2005, 2010 and 2011. The authors state that “to maintain consistent counts the majority of observations were made by the first author; a few observations were performed by three others in the vehicle with the first author and 12% of the observations were done by the second author after extensive travel with the first author”.

RESOURCES REQUIRED

In general, trained staff are the key resource needs to undertake the surveys on foot, boat and by vehicle. The additional costs of vehicles, including boats and helicopters may also need to be considered depending upon the geography, habitats and size of the areas invaded. If remote sensing is being adopted then drones and computer software are also needed, along with staff with the relevant skills.

For the digital aerial sketch-mapping system (DASM) detailed above, Rebbeck *et al.*, (2015) report for the USA that approximately 12,000 acres were surveyed per day at an average of 2,000 acres/hour. They remark that there is a distinct cost advantage to their sketch-mapping technique over ground-based detection only. Their survey costs were estimated at \$0.40/acres (\$1/ha). Additional populations of non-seed-producing *Ailanthus* were often found during ground searches of seed-bearing *Ailanthus* for chemical treatments. This suggests that a multistage approach of conducting local ground searches near aerially detected seed-bearing *Ailanthus* may be even more cost-effective. Rebbeck *et al.*, (2015) based the cost estimation on a commercial rate of \$960/hour, which included helicopter rental (\$800/hour) and hourly wages for the pilot (\$60/hour) and two sketch-mappers (\$100/hour) for an average 8-hour day. This estimate included round-trip flight time to and from the airport and survey area (1–2 hours/day), as well as a midday aircraft refuelling and crew break at the closest airport. Additional costs for training, setup, and post data processing time were minimal since they retooled an existing methodology currently used by the Division of Forestry. However, they estimated that an additional 16 hours (\$50/hour) is required annually. Estimates of coverage ranged from 2,000 to 7,000 acres/hour, depending on the level of *Ailanthus* infestation and flight altitude. If more *Ailanthus* was present, then more time per acre was required for mapping (McAvoy *et al.*, 2012).

SIDE EFFECTS

There will be no side effects in relation to early detection measures applied to tackle *Ailanthus*. However, if there were common biosecurity strategies for a number of invasive alien plants this will of course produce general beneficial effects as some vectors and corridors are responsible for the spread of more than one taxon, so that land surveillance in the same localities or along the same routes will tackle more than one alien taxon.

ACCEPTABILITY TO STAKEHOLDERS

See the sections on Preventive measures for unintentional introduction and spread. As generally agreed, public participation in detecting invasive species can increase the available “eyes and ears” searching for identified targets. A potential area of conflict is the access to private land to undertake the survey work.

ADDITIONAL COST INFORMATION

As *Ailanthus* is already present in many EU Member States, a lack of surveillance would certainly promote further spread. Early detection and rapid eradication of incipient populations of *Ailanthus* before they have a chance to become widely established, will eliminate the need for costly and resource-intensive control programs.

LEVEL OF CONFIDENCE*

High.

There is plenty of literature and practical cases supporting the fact that Early Detection, followed by Rapid Eradication (EDRE) would be a very effective strategy to limit further spread of *Ailanthus* within the EU. However, there is not enough information to calculate the total cost for the EU for such a strategic option.

Several measures do exist and would be very effective if included in a dedicated plan. However, such a dedicated plan should be based also on the knowledge of the actual distribution and abundance at MS level, at least with the resolution of a 10 × 10 km grid map (or even lower for some priority sites). Such important baseline mapping dataset is presently not available, so that the precise evaluation of efforts and resources required for ED in areas not yet invaded by *Ailanthus* in the EU is not possible.

Ailanthus altissima. © Wolfgang Rabitsch



* See Appendix

Measures to achieve rapid eradication after an early detection of a new occurrence.



Hand pulling.

MEASURE DESCRIPTION

Early Detection, followed by Rapid Eradication (EDRE), can detect and eradicate incipient populations of *Ailanthus* before they have a chance to become widely established, thus eliminating the need for costly and resource-intensive control programs. If prevention fails, early detection and rapid response are the next and most cost-effective line of defence against invasive alien species.

The measures to achieve rapid eradication of *Ailanthus* are the same as described in the section on Management, for example, rapid eradication should follow an integrated control methodology.

In the very first phase of an invasion where only seedlings are present (before the tap root develops) hand pulling can be applied in combination with monitoring of the site and control follow ups. In the case of larger infestations, vegetative propagation from adult individuals, rapid eradication should be conducted according to an integrated control methodology in the framework of a management plan (see *Management measures* sections below). If rapid eradication occurs on a relatively large area (such as more than 10 m²), it is advisable to adopt specific measures for vegetation recovering of the eradicated areas. These might include the planting or sowing of local plant species or the temporary protection from grazing. For these reasons, rapid eradication cannot be applied without considering an integrated control methodology.

EFFECTIVENESS OF MEASURE

Rapid eradication is expected to be very effective. Rejmanek and Pitcairn (2002) report some of the numerous examples where small infestations of invasive plant species have been eradicated by hand pulling. According to the study the same authors conducted in California, professional eradication of exotic weed infestations smaller than one hectare are usually possible.

Efforts in eradicating first establishment stages of *Ailanthus* are likely more challenging than in many other species as young plants, rather quickly, develop a tap-root and respond

with vegetative reproduction to disturbance. Thus, control for root suckers may be vital for a period of time longer than 2 years.

EFFORT REQUIRED

Manpower and simple tools. As a result of the presence of an *Ailanthus* seed bank, the removal by hand may need to be repeated several times in a 2-year period. Seedlings will need to be monitored and removed thereafter (by hand pulling). According to Rebeck *et al.*, (2010) persistence of *Ailanthus* seeds in soil seed banks is underreported. It appears that seeds are short-lived, typically 1 to 2 years, but germination rates are very high (80 to 100 percent) in disturbed forest stands.

RESOURCES REQUIRED

The resources required will be related to the extent of the invaded area, the land accessibility, the need to repeat the interventions. When *Ailanthus* hand pulling is planned, it is important to use gloves to prevent skin irritation. DGE-BIODIV (2013)⁵ give costs in Switzerland for manual pulling at 3–7 CHF/m²/year.

SIDE EFFECTS

Eradication actions need to be planned carefully and restoration actions are always necessary as disturbance during interventions might promote invasion by other generalist invasive alien plants and *Ailanthus* vegetative spread. Restoration of undisturbed native vegetation might be a way to limit further spread of *Ailanthus*. Hand pulling of young seedlings and suckers should be avoided (or carefully performed) from historical (or archaeological) buildings to prevent further damages on the structures (Vidotto, 2015).

ACCEPTABILITY TO STAKEHOLDERS

See the sections on preventive measures for unintentional introduction and spread.

Public perception of localised control of *Ailanthus* seedlings and young plants is generally positive when found outside cultivation and in land-uses where it is considered a weed. On the contrary, the removal of large trees can be conflictual

⁵ http://www.vd.ch/fileadmin/user_upload/themes/environnement/faune_nature/fichiers_pdf/Flore_et_Champignons/Recommandations_de_lutte_VD_F4-1_-_Ailante.pdf

in some cases. To promote acceptability, it can be useful to identify and promote replacement plant species for *Ailanthus* after its eradication from private or public gardens and urban green areas.


ADDITIONAL COST INFORMATION

As *Ailanthus* is already present in many EU Member States, a lack of intervention, in particular local eradication of new outbreaks, would certainly promote a further spread. Eradication of incipient populations of invasive plant species before they have a chance to become widely established, will eliminate the need for costly and resource-intensive control programs. In addition, it is generally agreed that “doing nothing” or waiting for more research violates precautionary principles (Underwood, 1997). Thus, local authorities and land managers should plan programs to detect and facilitate the eradication of small, incipient *Ailanthus* populations to attempt, in accordance with precautionary principles, to preclude additional widespread infestations.

LEVEL OF CONFIDENCE*

High.

There is plenty of literature and practical cases supporting the fact that Early Detection, followed by Rapid Eradication (EDRE) would be a very effective strategy to limit further spread of *Ailanthus* within the EU.



Ailanthus altissima. © Tim Adriaens, INBO

* See Appendix

Measures for the species' management.



Integrated control strategy.

MEASURE DESCRIPTION

Different measures may be required at an individual site, and management should be frequently site-specific.

Conceptually, the management of *Ailanthus* needs to include a range of technologies and tools rather than only plant protection products (herbicides) and/or mechanical interventions alone. Different types of habitats and land uses are invaded by *Ailanthus* in the European Union, and within a single country or region. The idea of an **integrated control** originates from the agricultural sector but can be very effectively applied to many of the invasive alien plants that impact the environment. This is often referred as IPM 'integrated pest management', IWM 'integrated weed management', IVM 'integrated vegetation management' or also as best management practices (BMPs). Numerous reviews describing IWM can be referenced for specific details (Thill *et al.*, 1991; van Wilgen *et al.*, 2001; Owen *et al.*, 2015). The key principle to an IVM program is to preserve as much desirable vegetation as practical while removing the undesirable vegetation and the Pennsylvania Department of Transportation have proposed one for managing *Ailanthus* on roadsides (Gover *et al.*, 2004). Integrated control has been suggested also for Switzerland⁶, Hungary and Slovakia⁷.

In the case of *Ailanthus* **integrated control**, specific tactics can include any combination of the following: herbicide rotation and herbicide mixtures, herbicide application rate and timing, precision application (such as "drill-fill" or stem injection), selection and integration of different methods according to the location and site accessibility, age, size, and the sex of the targeted *Ailanthus* plant (hand pulling can be an option, even if in very few cases), seed bank management, habitat manipulation (both for prevention and control) which can include several management and silvicultural options and disturbance mitigation. Importantly, as studied in the US, even rodents can have a significant effect on the survival of tree seeds and seedlings invading old fields (Ostfeld *et al.*, 1997).

Mechanical and chemical control are short-term activities, whereas rigorous and disciplined follow-up and rehabilitation are usually necessary in the medium term (van Wilgen *et al.*,

2001). In addition, control interventions are often likely to be upset by stochastic events such as fires, floods, annual climate variation on control outcomes or other events. While an approach of **adaptive management**, based on trial, error and continual improvement is a logical way in which to progress (van Wilgen *et al.*, 2001; Kettenring and Adams, 2011; Williams and Brown, 2016), integrated control should be always included in a dedicated **management plan** (site or habitat specific) to cope with stakeholders' issues, the need of monitoring, control follow ups, and prevention of new invasions (by other taxa) or re-invasion (by *Ailanthus*). Useful information can be achieved in the work of Leffler and Sheley (2012) describing how adaptive management could be included in an ecologically based invasive plant management strategy (EBIPM). EU/national/local legislation on the use of plant protection products and biocides needs to be always respected.

In many types of invaded sites mechanical and chemical control for *Ailanthus* are more effective when used in combination, therefore the two methods are often combined and are here treated as a single management measure. *Ailanthus* is very difficult or impossible to control by simple cutting or by other mechanical means alone. Not only do trees re-sprout with tremendous vigour, but massive root suckering also occurs, which usually results in many more new stems spreading over a wider area (USDA, 2014). There are several practical techniques that can help make **mechanical control** of tree-of-heaven more effective. These include: (1) cutting trees before they become too large; (2) cutting trees in early summer when root reserves are lowest; (3) cutting regrowth repeatedly and frequently, and applying Plant Protection Products to cut surfaces; and (4) providing shade from competitive native plants after control efforts (USDA, 2014). After larger trees have been cut to ground level, re-sprouting wood is soft enough to be mowed at regular intervals which can stress root reserves and lead to fewer root suckers. However, an infrequent mowing cycle may allow the sprouts to spread and become too tall to mow. It is important to stress that using mechanical treatment alone risks the measure being ineffective, as if one sprout survives it can be the source for establishing abundant populations again. Therefore, mowing – similarly to the other

6 http://www4.ti.ch/fileadmin/GENERALE/organismi/rapporti_studi/Ailanto_Oikos_2008.pdf

7 <https://www.dunaipoly.hu/uploads/2016-02/20160202200313-rosalia-handbook-ver2-6xtoafsq.pdf>

mechanical methods – is more effective when followed up with a chemical treatment (USDA, 2014).

Plant protection products (PPP) can be applied in a variety of ways including: (1) foliage application; (2) topical application to cut stems and stumps; (3) injection into the trunk; and (4) basal spraying. Although aboveground portions of *Ailanthus* are relatively easy to suppress or kill with herbicide treatment, it is also important to control the root system. Therefore, special attention should be paid to selection of the correct PPP, optimal application rate, and appropriate time to get good results. Care should always be taken when spraying any PPP near non-target plants. It is important to read and carefully follow all instructions and warnings provided on the PPP label (USDA, 2014). To reduce risk related to the use of PPP, the main principle of **targeted chemical control methods (TCCM)** should be followed. This means to apply the herbicides directly, and only, on specific plant individuals selected for control. TCCM were originally developed in order to remove specific invasive alien plant taxa from natural and protected areas without harming native species and sensitive habitats (Tu *et al.*, 2001; Burn, 2003). Targeted chemical control methods suggested for *Ailanthus* in Cyprus include 3 different techniques: drill-fill (stem injection), cut-stump, and frilling (Dufour-Dror, 2013). Drill-fill (stem injection) is considered the method that gives the best results for invasive tree such *Ailanthus* characterized by a high resilience and re-sprouting ability. Because drilling represents a localized wound in the tree structure, it is less likely to trigger intense root sucker's development as that happens when the main trunk is cut down. The cut-stump is the second type of TCCM widely used in order to control alien invasive trees. With the cut-stump technique the trunks are cut down prior to herbicide application (herbicide is applied directly to the cambial regions of the newly cut stem). Herbicides are not injected into the sapwood as it is with drill-fill but applied onto the stump (Dufour-Dror, 2013). Frilling is a TCCM suited for young saplings and seedlings (BD<5cm) that are too small to be drilled, and whose exposed cut surface is too limited in order to apply enough herbicide (Burn, 2003; Dufour-Dror, 2013). Drill-fill techniques were tested also in Island of Sicily (Italy) as described by Badalamenti and La Mantia (2013) and on the island of Montecristo (Italy).

Integrated control strategy within a dedicated management plan for *Ailanthus* can also include **grazing**. *Ailanthus* usually has low palatability for grazers; however, livestock (cattle, sheep, and goats) and deer will consume it during certain times of the year under specific circumstances. Goats will eat leaves and bark. Deer will browse leaves during the summer – especially in shady, forested areas. Under heavy grazing pressure, livestock will remove new suckers and sprouts. Use of grazing in combination with other techniques may be effective in certain situations

(USDA, 2014). However, grazing alone is unlikely to control large infestations. In addition, grazing cannot be proposed in some types of infested land uses, such as road verges or urban areas. Furthermore, grazing has to be completely avoided from sites where vegetation recovering is planned after a removal/eradication intervention against *Ailanthus*.

EFFECTIVENESS OF MEASURE

The control of *Ailanthus* by integrated control strategy within a dedicated management plan is an effective measure. Examples of effective control are reported by Brunel *et al.*, (2013).

On the contrary, single and non-integrated measures alone both mechanical (such as sapling removal or cutting) and chemical (such as foliar sprays or basal bark application) have met with limited success and are costly, time-consuming, and can pose a risk of serious non-target effects (Swearingen and Pannill, 2009; Erdős *et al.*, 2015).

The LIFE project Montecristo 2010 (Montecristo 2010: eradication of invasive plant and animal aliens and conservation of species/habitats in the Tuscan Archipelago, Italy, LIFE08 NAT/IT/000353) aimed to eradicate *Ailanthus* from the Italian islands of Montecristo and Pianosa (about 10 km²). The topography of Montecristo makes this intervention very difficult and costly, while on Pianosa the topography of the island is less a problem. So far, the abundance of *Ailanthus* has been significantly reduced on both islands, mostly on Pianosa and activities will continue until the full eradication of *Ailanthus* from both islands (http://www.montecristo2010.it/stealthV3_publica/0840425A051345033092.pdf).

EFFORT REQUIRED

The measures need to be maintained indefinitely.

RESOURCES REQUIRED

The resources needed for an integrated control strategy within a dedicated management plan will be dependent upon specific circumstances, but in general will include trained staff, access to necessary tools and equipment to undertake physical control, the purchasing and application of PPP, and the disposal costs for the removed plants/trees. See DGE-BIODIV (2013) for details of the recommended resources required for the different individual control methods.

There is only little information on the cost for applying an integrated control strategy within a dedicated management plan for *Ailanthus*. For example, for the control of *Ailanthus* scattered on a surface of about 180 ha on the Island of Montecristo (Italy) the dedicated LIFE Project (LIFE08 NAT/IT/000353)⁸ allocated a budget of about 380,000 Euro, 19,000 Euro for the action plan, and an additional budget of

8 http://www.montecristo2010.it/stealthV3_publica/0810372A007595000015.pdf

about 40,000 Euro for mapping *Ailanthus* distribution from helicopter and land survey. Management itself can incur direct, indirect, and opportunity costs where opportunity costs are the values that are forgone by implementing a particular management strategy (such as lost trade values from implementing a ban, or forgone profits from delaying timber harvest in infested woods). While direct expenditures on invasive species management are often straightforward to quantify and readily considered, opportunity costs are more often overlooked (Epanchin-Niell, 2017).

However, there are example of costs for single actions of an integrated control strategy, for example, for Switzerland, Germany, Italy (DGE-BIODIV, 2013 – Ailante. *Ailanthus altissima* (Mill.) Swingle. F4-1 Recommandations de lutte⁹). DGE-BIODIV (2013) provide annual costs in Switzerland for different control methods:

- Manual pulling, once per year = 3–7 CHF/m²/year
- Repeated mowing, 5–6 times per year = 1.5–3 CHF/m²/year
- Chemical treatment on leaves and stem, once per year = 1–5 CHF/m²/year
- Removing the plant/tree along with root system, once per year = 40–80 CHF/m²/year
- Felling of shrubs and trees and chemical treatment of the stumps, once per year = 15–60 CHF/m²/year
- Ring-barking, once per year = 15–30 CHF/m²/year
- Chemical impregnation of the bark, once per year = 6–15 CHF/m²/year
- Notching of the bark followed by chemical treatment, once per year = 25–30 CHF/m²/year
- Felling followed by repeated mowing, 5–6 times per year = 15–60 CHF/m²/year
- Felling followed by chemical treatment on leaves and stem, once per year = 15–60 CHF/m²/year.

SIDE EFFECTS

Ailanthus can have negative effects on human infrastructures and archaeological and historical buildings, so that reducing its abundance with management is likely to have a positive effect on the local economy. The control of *Ailanthus* with an integrated control strategy within a dedicated management plan will minimise all possible negative effects of mechanical treatment of *Ailanthus* alone (expensive and often not successful) or due to the use of PPP (herbicides) alone.

ACCEPTABILITY TO STAKEHOLDERS

See the sections on general considerations on the preventive measures for unintentional introduction and spread.

The successful implementation of any management plan and action is dependent, at least in part, on its acceptability to a wide range of stakeholders. Management actions can be accepted by stakeholders if well communicated, however this cannot be easily predicted as there are

always very diverse perceptions among stakeholders' groups, with special concern to alien trees that can have multiple uses, or cultural or ornamental value. However, the Spanish environmental managers surveyed by Andreu *et al.*, (2009) identified the following taxa as noxious: *Carpobrotus* spp., *Eucalyptus* spp., *Ailanthus altissima*, and *Robinia pseudoacacia*.

In regards to the application of PPP's it is important to note that EU/national/local legislation on the use of plant protection products and biocides needs to be respected.

Importantly, the European Commission has decided on the 10th January 2017 to register a European Citizens Initiative (ECI) inviting the Commission "to propose to Member States a ban on glyphosate, to reform the pesticide approval procedure, and to set EU-wide mandatory reduction targets for pesticide use" (http://europa.eu/rapid/press-release_IP-17-28_en.htm).

In Italy, the National Action Plan for the sustainable use of plant protection products (D.M. January 22, 2014 - *Decreto 22 gennaio 2014, Piano di azione nazionale per l'uso sostenibile dei prodotti fitosanitari*) is leading to limitations in the use of chemical pesticides on roads (Action A.5.5) and in population centres (Action A.5.6) (Lorenzini, 2016), for example, in land use types very commonly invaded by *Ailanthus*.

Very recently, Kissane and Shephard (2017) have examined the published literature to evaluate the extent to which glyphosate based herbicides can cause toxic effects in wildlife and people and the implications of chronic exposure. The main principle of targeted chemical control methods (TCCM) is to apply the herbicides directly, and only, on specific plant individuals selected for control are therefore always suggested whenever the use of PPP is planned.

ADDITIONAL COST INFORMATION

No information available.

However, it should be taken into consideration that additional cost could be required for the management of those plants of *Ailanthus* that have been intentionally planted as single plants or in plantations, and that may have a particular historical, cultural, economic or ornamental value, such as monumental trees (Zapponi *et al.*, 2017) or tree individuals introduced for the first time in a Botanic Garden.

LEVEL OF CONFIDENCE*

High.

There is significant scientific literature and practical evidence to support the content of this section with concern to the types of measure that can be applied and their

9 http://www.vd.ch/fileadmin/user_upload/themes/environnement/faune_nature/fichiers_pdf/Flore_et_Champignons/Recommandations_de_lutte_VD_F4-1_-_Ailante.pdf

* See Appendix

effectiveness. Several management options do exist and would be very effective if included in a dedicated strategic management plan. However, such a dedicated management plan should be based on the knowledge of the actual distribution and abundance at MS level, at least with the

resolution of a 10 × 10 km grid map (or even lower for some priority sites). Such important baseline mapping dataset is presently not available, so that the precise evaluation of efforts and resources required for managing *Ailanthus* in the EU is not possible.



Classical biological control.

MEASURE DESCRIPTION

A variety of insects and diseases affect *Ailanthus* with varying success; however, there are no biological control agents currently approved for use in the United States (USDA, 2014). In addition, it should be borne in mind that the release of macro-organisms as biological control agents is currently not regulated at EU level. Nevertheless national/regional laws are to be respected. Before any release of an alien species as a biological control agent an appropriate risk assessment should be made.

Among biological agents currently being researched in the USA, there are three fungal pathogens, two weevils, and a moth. Approval for one of the weevil species, *Eucryptorrhynchus brandti* (Coleoptera: Curculionidae), is

anticipated in the near future (Ding *et al.*, 2006; USDA, 2014).

Importantly, the biological control of *Ailanthus* using the wilt-inducing fungi in the genus *Verticillium* spp. as “mycoherbicides” could be a promising alternative in Europe (Maschek and Halmschlager, 2017).

A strain of *Verticillium albo-atrum* found in southern Pennsylvania by Don Davis and colleagues has been found to cause near complete mortality of *Ailanthus* in artificially infected stands (Harris *et al.*, 2013). However, recent phylogenetic and taxonomic studies revealed five new *Verticillium* species. Two of these, *V. alfalfae* Inderb. *et al.*, and *V. nonalfalfae* Inderb. *et al.*, resembled the

Ailanthus altissima. © Jean-Marc Dufour-Dror



distantly related *V. albo-atrum* s.s. in morphology. Thus, all prior morphology-based identifications and assigned host ranges of *V. albo-atrum* s.l. are to be questioned. Currently, only *V. nonalfalfae* and *V. dahliae* have been reported from wilting *Ailanthus* in Europe (Maschek and Halmschlager, 2017). In Europe, an Austrian isolate of *V. nonalfalfae*, which is considered the most effective *Verticillium* species as biocontrol agent against *Ailanthus*, was tested in field inoculations studies. Results revealed that disease progression on *Ailanthus* is negatively influenced by high temperatures in summer. Therefore, inoculation in spring turned out to be the most effective to combat *Ailanthus*. However, due to the common occurrence of *V. dahliae* in Austria, this pathogen might be considered as alternative biocontrol agent of *Ailanthus* on sites in warmer climates. For that purpose, further investigations of the host range and non-target effects of selected *V. dahliae* strains should be carried out according to Maschek and Halmschlager (2017).

EFFECTIVENESS OF MEASURE

The biological control of *Ailanthus* using the wilt-inducing fungi in the genus *Verticillium* spp. as “mycoherbicides” could be a promising alternative to other control measures (Maschek and Halmschlager, 2017). The huge potential of selected strains of *V. nonalfalfae* as biocontrol agents against *Ailanthus* already has been demonstrated in the United States (Maschek and Halmschlager, 2017 and references cited therein) and Europe. *Verticillium dahliae* might serve as alternative biocontrol agent of *Ailanthus* on sites in warmer climates, however, further investigations

concerning host range and non-target effects of selected *V. dahliae* strains have to be carried out (Maschek and Halmschlager, 2017).

EFFORT REQUIRED

No information available.

RESOURCES REQUIRED

No information available.

SIDE EFFECTS

The application of a pathogen such as a *Verticillium* species bears the risk of non-target effects, particularly if it has a broad host range.

ACCEPTABILITY TO STAKEHOLDERS

The release of macro-organisms as biological control agents is currently not regulated at EU level. Nevertheless national/regional laws are to be respected. Before any release of an alien species as a biological control agent an appropriate risk assessment should be made.

ADDITIONAL COST INFORMATION

No information available.

LEVEL OF CONFIDENCE*

High.

Classical biological control could be a very useful methodology for long term management of invasive weeds.

* See Appendix

Bibliography

- Andreu, J., Vilà, M. and Hulme, P.E. (2009). An assessment of stakeholder perceptions and management of noxious alien plants in Spain. *Environmental Management*, 43: 1244–1255, DOI 10.1007/s00267-009-9280-1.
- Badalamenti, E. and La Mantia, T. (2013). Stem-injection of herbicide for control of *Ailanthus altissima* (Mill.) Swingle: a practical source of power for drilling holes in stems. *iForest* 6: 123–126 [online 2013-03-05] URL: <http://www.sisef.it/forest/contents?id=ifor0693-006>
- Ballero, M., Ariu, A. and Falagiani Piu, P.G. (2003). Allergy to *Ailanthus altissima* (tree of heaven) pollen. *Allergy*, 58: 532–533. doi: 10.1034/j.1398-9995.2003.00172.x
- Baptista, P., Costa, A.P., Simões, R. and Amaral, M.E. (2014) *Ailanthus altissima*: An alternative fiber source for papermaking. *Industrial Crops and Products*, 52: 32– 37.
- Boer, E. (2013). Risk assessment *Ailanthus altissima* (Mill.) Swingle. Naturalis Biodiversity Center, Leiden (<https://www.nvwa.nl/>).
- Brunel, S., Brundu, G. and Fried, G. (2013). Eradication and control of invasive alien plants in the Mediterranean Basin: towards better coordination to enhance existing initiatives. *Bulletin OEPP/EPPO Bulletin*, 43 (2): 290–308. DOI: 10.1111/epp.12041.
- Burn, A. (2003). The herbicide handbook: Guidance on the use of herbicides on nature conservation sites. English Nature in association with FACT. Peterborough, UK. p.195.
- Cabra-Rivas, I., Alonso, A. and Castro-Díez, P. (2014). Does stream structure affect dispersal by water? A case study of the invasive tree *Ailanthus altissima* in Spain. *Management of Biological Invasions*, 5 (2): 179–186, DOI: <http://dx.doi.org/10.3391/mbi.2014.5.2.11>.
- Casella, F. and Vurro, M. (2012). Mappatura di un'infestante arborea in aree extra-agricole: il caso dell'ailanto in Puglia. *ATTI Giornate Fitopatologiche*, 2012, 1, 695-704.
- Celesti-Grapow, L. and Blasi, C. (2004). The role of alien and native weeds in the deterioration of archaeological remains in Italy. *Weed Technology*, 18, Invasive Weed Symposium: 1508-1513.
- Conedera, M., Nobis, M. and Wunder, J. (2014). Ailanto: Ospite indesiderato in bosco? *Sherwood*, 206: 15-18.
- Ding, J., Reardon, R., Wu, Y., Zheng, H. and Fu, W. (2006). Biological control of invasive plants through collaboration between China and the United States of America: a perspective. *Biological Invasions*, 8:1439–1450, DOI 10.1007/s10530-005-5833-2
- Dufour-Dror, J-M. (2013). Guide for the control of invasive trees in natural areas in Cyprus: strategies and technical aspects. Department of Forests, Republic of Cyprus, 25 pp. (http://www.amiandos.eu/images/docs/Guide_for_the_Control_of_Invasive_Trees_in_Natural_Areas_in_Cyprus_FINAL_16-12-2013.pdf).
- Epanchin-Niell, R.S. (2017). Economics of invasive species policy and management. *Biological Invasions*, DOI 10.1007/s10530-017-1406-4.
- Erdős, L., Márkus, A. and Körmöczy, L. (2005). Consequences of an extirpation trial of the tree of heaven (*Ailanthus altissima* (Mill.) Swingle) on rock grasslands and slope steppes. *Tiscia*, 35: 3-7.
- Feret, P.P. (1985). *Ailanthus*: variation, cultivation, and frustration. *Arboric Urban For.*, 11 (12): 361–368.
- Fryer, J. L. (2010). *Ailanthus altissima*. In: *Fire Effects Information System*, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (<http://www.fs.fed.us/database/feis/>).
- Gover, A., Kuhns, L. and Johnson, J. (2004). Managing tree-of-Heaven (*Ailanthus altissima*) on Roadsides. Roadside Vegetation Management, Pennsylvania Department of Transportation, Factsheet 3, 4 pp. (<http://plantscience.psu.edu/research/projects/vegetative-management/publications/roadside-vegetative-mangement-factsheets/3ailanthus-on-roadsides>).
- Graziano, G. and Clayton, L. (2017). Best management practices. Controlling the spread of invasive plants during road maintenance. University of Alaska Fairbanks, 12-14/GG-SS-LC/2-17, 39 pp. (<https://www.uaf.edu/files/ces/publications-db/catalog/anr/PMC-00342.pdf>).
- Halford, M., Heemers, L., van Wesemael, D., Mathys, C., Wallens, S., Branquart, E., Vanderhoeven, S., Monty, A. and Mahy, G. (2014). The voluntary Code of conduct on invasive alien plants in Belgium: results and lessons learned from the AlterIAS LIFE+ project. *EPPO Bulletin*, 44: 212–222. DOI: 10.1111/epp.12111.
- Harris, P.T., Cannon, G.H., Smith, N.E. and Muth, N.Z. (2013). Assessment of plant community restoration following tree-of-Heaven (*Ailanthus altissima*) control by *Verticillium albo-atrum*. *Biological Invasions*, 15:1887–1893, DOI: 10.1007/s10530-013-0430-2
- Harris, S., Brown, J. and Timmins, S. (2001). Weed surveillance – how often to search. *Science for Conservation*, 175, 27 pp. (<http://www.doc.govt.nz/documents/science-and-technical/SFC175.pdf>).
- Holec, J., Vodičková, E. and Soukup, J. (2014). Sex composition of *Ailanthus altissima* population in the city of Prague and the occurrence of andromonoecious individuals. In Book of Abstract, 8th International Conference on Biological Invasions from understanding to action, p. 228. (http://www.esenias.org/files/NEOBIOA_2014_Proceedings.pdf).
- Kettenring, K.M. and Adams, C.R. (2011). Lessons learned from invasive plant control experiments: a systematic review and meta-analysis. *Journal of Applied Ecology*, 48: 970–979. doi: 10.1111/j.1365-2664.2011.01979.x.
- Kissane, Z. and Shephard, J.M. (2017). The rise of glyphosate and new opportunities for biosentinel early-warning studies. *Conservation Biology*, DOI: 10.1111/cobi.12955.
- Kota, N.L., Landenberger, R.E. and McGrawm G.B. (2007). Germination and early growth of *Ailanthus* and tulip poplar in three levels of forest disturbance. *Biological Invasions*, 9: 197–211, DOI 10.1007/s10530-006-9026-4.
- Kowarik, I. (1983). Zur Einbürgerung und zum pflanzengeographischen Verhalten des Götterbaumes (*Ailanthus altissima* (Mill.) Swingle) im französischen Mittelmeergebiet (Bas-Languedoc). *Phytocoenologia*, 11: 389–405.
- Kowarik, I. and Säumel, I. (2007). Biological flora of Central Europe: *A. altissima* (Mill.) Swingle Perspect. *Plant Ecol Evol Syst*, 8: 207–237.
- Kowarik, I. and Säumel, I. (2008). Water dispersal as an additional pathway to invasions by the primarily wind-dispersed tree *Ailanthus altissima*. *Plant Ecology*, 198: 241–252, DOI 10.1007/s11258-008-9398-x.
- Kowarik, I. and von der Lippe, M. (2011). Secondary wind dispersal enhances long-distance dispersal of an invasive species in urban road corridors. *NeoBiota*, 9: 49-70.
- Knüsel, S., Conedera, M., Rigling, A., Fonti, P. and Wunder, J. (2015). A tree-ring perspective on the invasion of *Ailanthus altissima* in protection forests. *Forest Ecology and Management*. <http://dx.doi.org/10.1016/j.foreco.2015.05.010>.
- Landenberger, R.E., Kota, N.L. and McGraw, J.B. (2007). Seed dispersal of the non-native invasive tree *Ailanthus altissima* into contrasting environments. *Plant Ecology*, 192: 55–70. <https://doi.org/10.1007/s11258-006-9226-0>.

- Leffler, A.J. and Sheley, R.L. (2012). Adaptive management in EBIPM. *Rangelands*, 34(6): 44–47. <https://doi.org/10.2111/RANGELANDS-D-12-00053.1>
- Lorenzini, G. (2016). Will a fungus save us from the *Ailanthus* invasion? *Italian Journal of Mycology*, [S.l.], v. 45, p. 13-18, may 2016. ISSN 2531-7342 (<https://italianmycology.unibo.it/article/view/6151>).
- Marco, A., Lavergne, S., Dutoit, T. and Bertaudiere-Montes, V. (2010). From the backyard to the backcountry: how ecological and biological traits explain the escape of garden plants into Mediterranean old fields. *Biological Invasions*, 12: 761–779. DOI 10.1007/s10530-009-9479-3.
- Maschek, O. and Halmschlager, E. (2017). Natural distribution of *Verticillium* wilt on invasive *Ailanthus altissima* in eastern Austria and its potential for biocontrol. *Forest Pathology*. e12356. <https://doi.org/10.1111/efp.12356>
- McAvoy, T.J., Snyder, A.L., Johnson, N., Salom, S.M. and Kok, L.T. (2012). Road Survey of the Invasive tree-of-Heaven (*Ailanthus altissima*) in Virginia. *Invasive Plant Science and Management*, 5(4): 506– 512, <https://doi.org/10.1614/IPSM-D-12-00039.1>.
- Merriam, R.W. (2003). The abundance, distribution and edge associations of six non-indigenous, harmful plants across North Carolina. *J. Torrey Bot. Soc.*, 130: 283–291.
- Murina, M. and Nicita, A. (2017). Trading with conditions: The effect of sanitary and phytosanitary measures on the agricultural exports from low-income countries. *The World Economy*, DOI: 10.1111/twec.12368.
- Napoli, M. (2008). The plants, rituals and spells that ‘cured’ helminthiasis in Sicily. *Journal of Ethnobiology and Ethnomedicine* 4:21, doi:10.1186/1746-4269-4-21.
- Ostfeld, R.S., Manson, R.H. and Canham, C.D. (1997). Effects of rodents on survival of tree seeds and seedlings invading old fields. *Ecology*, 78(5): 1531–1542.
- Owen, M.D.K., Beckie, H.J., Leeson, J.Y., Norsworthy, J.K. and Steckel, L.E. (2015). Integrated pest management and weed management in the United States and Canada. *Pest Manag Sci*, 71: 357–376.
- Peng, S., Yi, Z. and Liu, M. (2017) Ailanthone: a new potential drug for castration-resistant prostate cancer. *J Cancer*, 36:25. DOI 10.1186/s40880-017-0194-7
- Pennsylvania Department of Transportation. (2014). Invasive Species Best Management Practices. Pub 756 (4/14), 43 pp. (<http://www.dot.state.pa.us/public/pubsforms/Publications/PUB%20756.pdf>).
- Radtke, A., Ambrass, S., Zerbe, S., Tonon, G., Fontana, V. and Ammer, C. (2013). Traditional coppice forest management drives the invasion of *Ailanthus altissima* and *Robinia pseudoacacia* into deciduous forests. *For Ecol Manag*, 291:308–317.
- Rebbeck, J., Hutchinson, T.F., Iverson, L., Yaussy, D., Boyles, R. and Bowden, M. (2010). Studying the effects of management practices on *Ailanthus* populations in Ohio forests: a research update. USDA Research Forum on Invasive Species, GTR-NRS-P-75, 50-51 (<https://www.nrs.fs.fed.us/pubs/gtr/gtr-nrs-p-75papers/24rebbeck-p-75.pdf>).
- Rebbeck, J., Kloss, A., Bowden, M., Coon, C., Hutchinson, T.F., Iverson, L. and Guess, G. (2015). Aerial Detection of Seed-Bearing Female *Ailanthus altissima*: A cost-effective method to map an invasive tree in forested landscapes. *Forest Science*, 61(6): 1068–1078, <http://dx.doi.org/10.5849/forsci.14-223>.
- Rejmánek, M. and Pitcairn, M.J. (2002). When is eradication of exotic pest plants a realistic goal? In: Turning the tide: the eradication of invasive species, C.R. Veitch and M.N. Clout (eds.), pp. 249–53. IUCN SSC Invasive Species Specialist Group, Gland, Switzerland, and Cambridge, UK.
- Säumel, I. and Kowarik, I. (2010). Urban rivers as dispersal corridors for primarily wind-dispersed invasive tree species. *Landscape and Urban Planning*, 94: 244–249.
- Swearingen, J. and Pannill, P. (2009). Least wanted: *Ailanthus altissima*. Plant Conservation Alliance. <http://www.nps.gov/plants/alien/fact/aial1.htm>
- Thill, D.C., Lish, J.M., Callihan, R.H. and Bechinski, E.J. (1991). Integrated weed management: a component of integrated pest management: a critical review. *Weed Technology*, 5:648–656.
- Tsao R., Romanchuk F.E., Peterson C.J. and Coats J.R. (2002). Plant growth regulatory effect and insecticidal activity of the extracts of the tree of Heaven (*Ailanthus altissima* L.). *BMC Ecology*, 2:1, This article is available from: <http://www.biomedcentral.com/1472-6785/2/1>.
- Tu, M., Hurd, C., Robinso, R. and Randall, J.M. (2001). Weed control methods handbook. The Nature Conservancy. p. 218.
- Underwood, A.J. (1997). Environmental decision making and the precautionary principle: what does this principle mean in environmental sampling practice? *Landscape and Urban Planning*, 37: 137–146.
- USDA Forest Service San Dimas Technology. (2017). Dangerous travelers: Controlling invasive plants along America’s roadways. (<https://www.fs.fed.us/invasivespecies/prevention/dangeroustravelers.shtml>).
- USDA, United States Department of Agriculture. (2014). Field Guide for managing tree-of-heaven in the Southwest. TP-R3-16-09, 9 pp. (https://www.fs.usda.gov/Internet/FSE_DOCUMENTS/stelprdb5410131.pdf).
- van Wilgen, B., Richardson, D.M. and Higgins, S. (2001). Integrated control of invasive alien plants in terrestrial ecosystems. *Land Use and Water Resources Research*, 1 (5): 1–6.
- Vidotto, F., Fogliatto, S., Milan, M., Destefani, G., Gatti-Spriano, I. and Ferrero, A. (2015). The destructive invasion by *Ailanthus altissima* (Mill.) Swingle of the eighteenth-century fortress “Cittadella” of Alessandria. Book of Abstracts, 17th European Weed Research Society Symposium - EWRS 2015, p. 261 (<https://iris.unito.it/handle/2318/1521435?mode=full.277#WXopFOlLdPY>).
- von der Lippe, M., Bullock, J.M., Kowarik, I., Knopp, T. and Wichmann, M. (2013). Human-mediated dispersal of seeds by the airflow of vehicles. *PLOS ONE*, 8(1): e52733.
- Wickert, K.L., O’Neal, E.S., Davis, D.D. and Kasson, M.T. (2017). Seed production, viability, and reproductive limits of the invasive *Ailanthus altissima* (tree-of-Heaven) within invaded environments. *Forests*, 8, 226; doi:10.3390/f8070226.
- Williams, B.K. and Brown, E.D. (2016). Technical challenges in the application of adaptive management. *Biological Conservation*, 195: 255–263. <http://dx.doi.org/10.1016/j.biocon.2016.01.012>
- Wunder, J., Knüsel, S., Gurtner, D. and Conedera, M. (2016). 5.7 The spread of tree of heaven in Switzerland. In: Krumm, F. and Vitková, L. (eds) 2016. *Introduced tree species in European forests: opportunities and challenges*, pp. 374–385. European Forest Institute (http://www.in-tree.org/uploads/images/book/Introduced_tree_species_EN_HighRes.pdf).
- Zapponi, L., Mazza, G., Farina, A., Fedrigoli, L., Mazzocchi, F., Roversi, P.F., Sabbatini Peverieri, G. and Mason, F. (2017). The role of monumental trees for the preservation of saproxylic biodiversity: re-thinking their management in cultural landscapes. *Nature Conservation*, 19: 231–243, <https://doi.org/10.3897/natureconservation.19.12464>.

WEB sites

LIFE Montecristo

http://www.minambiente.it/sites/default/files/archivio/allegati/life/life_progetto_montecristo_action_plan.pdf

LIFE Alta Murgia

http://ec.europa.eu/environment/life/project/Projects/index.cfm?fuseaction=search.dspPageandn_proj_id=4566anddocType=pdf

LIFE RI.CO.PR.I

http://www.pdc.minambiente.it/sites/default/files/progetti/ricopri_linee-guida-azione-c2.pdf

Modelling *Ailanthus* distribution

<http://www.gissmox.ethz.ch/en/showcases/projects/ailanthus.html>

***Ailanthus* control methods**

Switzerland (protected area)

<http://www.gissmox.ethz.ch/en/showcases/projects/ailanthus.html>

Switzerland (forest)

<https://www.bafu.admin.ch/bafu/it/home/temi/bosco/pubblicazioni-studi/pubblicazioni/guida-relativa-alla-gestione-dell-ailanto.html>

http://www4.ti.ch/fileadmin/GENERALE/organismi/rapporti_studi/Ailanto_Oikos_2008.pdf

Italy (historical site, page 6 of the PDF document)

http://www.isaitalia.org/images/stories/arbore/rivista_elettronica/2016/2/Arbor_n_2_2016-ilovepdf-compressed1.pdf

https://iris.unito.it/retrieve/handle/2318/1589122/206577/Vidotto_etal_ailanto_arbor_postprint_4aperto.pdf

Italy

http://www.regione.piemonte.it/ambiente/tutela_amb/dwd/esotichelInvasive/schede_complete/ailanto_althissima_2016.pdf

France

http://www.codeplantesenvahissantes.fr/fileadmin/PEE_Ressources/TELECHARGEMENT/Ailanthus_altissima__Mill__Swingle.pdf

Spain

http://www.mapama.gob.es/es/biodiversidad/temas/conservacion-de-especies/ailanthus_altissima_2013_tcm7-306916.pdf

http://www.lifemedwetrivers.eu/sites/default/files/documentos/25_ailanthus_altissima.pdf

Appendix

- **High: Well established:** comprehensive meta-analysis or other synthesis or multiple independent studies that agree. Note: a meta-analysis is a statistical method for combining results from different studies which aims to identify patterns among study results, sources of disagreement among those results, or other relationships that may come to light in the context of multiple studies.
- **Established but incomplete:** general agreement although only a limited number of studies exist but no comprehensive synthesis and/or the studies that exist imprecisely address the question.
- **Unresolved:** multiple independent studies exist but conclusions do not agree.
- **Inconclusive:** limited evidence, recognising major knowledge gaps.

Your feedback is important. Any comments that could help improve this document can be sent to ENV-IAS@ec.europa.eu

This technical note has been drafted by a team of experts under the supervision of IUCN within the framework of the contract No 07.0202/2016/739524/SER/ENV.D.2 "Technical and Scientific support in relation to the Implementation of Regulation 1143/2014 on Invasive Alien Species". The information and views set out in this note do not necessarily reflect the official opinion of the Commission. The Commission does not guarantee the accuracy of the data included in this note. Neither the Commission nor any person acting on the Commission's behalf may be held responsible for the use which may be made of the information contained therein. Reproduction is authorised provided the source is acknowledged. Reuse is authorised provided the source is acknowledged. For any use or reproduction of photos or other material that is not under the EU copyright, permission must be sought directly from the copyright holders.