



18-23439 (17-23202)

Pest Risk Analysis for *Triadica sebifera*



2018

EPPO

21 Boulevard Richard Lenoir
75011 Paris
www.eppo.int hq@eppo.int

This pest risk analysis scheme has been specifically amended from the EPPO Decision-Support Scheme for an Express Pest Risk Analysis document PM 5/5(1) to incorporate the minimum requirements for risk assessment when considering invasive alien plant species under the EU Regulation 1143/2014. Amendments and use are specific to the LIFE Project (LIFE15 PRE FR 001) 'Mitigating the threat of invasive alien plants to the EU through pest risk analysis to support the Regulation 1143/2014'.

Cite this document as:

EPPO (2018) *Pest risk analysis for Triadica sebifera* EPPO, Paris.

Available at: XXX

Photo: *Triadica sebifera* infestation in North America (Image courtesy of Nancy Loewenstein, Auburn University, Bugwood.org)

EUROPEAN AND MEDITERRANEAN PLANT PROTECTION ORGANIZATION

Pest risk analysis for *Triadica sebifera*

This PRA follows EPPO Standard PM5/5 Decision support scheme for an Express Pest Risk Analysis

Composition of the Expert Working Group

BRUNDU Giuseppe (Mr)	University of Sassari, Department of Agriculture, Viale Italia 39, 07100 Sassari, Italy gbrundu@tin.it
Chapman Daniel (Mr)	Centre for Ecology and Hydrology Bush Estate, Penicuik, Edinburgh , UK, dcha@ceh.ac.uk
FLORY S. Luke (Mr)	Agronomy Department, University of Florida, PO Box 110500, FL 32611 Gainesville, United States flory@ufl.edu
LE ROUX Jaco (Mr)	Department of Botany and Zoology, Stellenbosh University, Private Bag X1, 7602 Matieland, South Africa jleroux@sun.ac.za
PESCOTT Oliver (Mr)	CEH Wallingford, Benson Lane, OX10 8BB Wallingford, Oxfordshire, United Kingdom olipes@nerc.ac.uk
SIEMANN Evan (Mr)	Rice University, 6100 S. Main St, TX 77005 Houston, United States siemann@rice.edu
STARFINGER Uwe (Mr)	Julius Kühn Institut (JKI), Federal Research Centre for Cultivated Plants, Institute for National and International Plant Health, Messeweg 11/12, 38104 Braunschweig, Germany uwe.starfinger@julius-kuehn.de
TANNER Rob (Mr)	OEPP/EPPO, 21 boulevard Richard Lenoir, 75011 Paris, France Tel: +33-145207794 - rt@epo.int

The pest risk analysis for *Triadica sebifera* has been performed under the LIFE funded project:



LIFE15 PRE FR 001

Mitigating the threat of invasive alien plants to the EU through pest risk analysis to support the Regulation 1143/2014

In partnership with

EUROPEAN AND MEDITERRANEAN PLANT PROTECTION ORGANIZATION

And

NERC CENTRE FOR ECOLOGY AND HYDROLOGY



Review Process

- This PRA on **Triadica sebifera** was first drafted by S. Luke Flory and Austin Young, University of Florida.
- The PRA was further evaluated by international experts which made up an Expert Working group that physically met in Paris in 2017.
- The PRA has been reviewed by the EPPO Panel on Invasive Alien Plants in 2017.
- The PRA was reviewed by the EPPO Core Members for PRA (2017/18)
- The PRA was reviewed by the EU Scientific Forum (2018)

Contents

Summary	6
Stage 1: Initiation	10
Stage 2: Pest Risk Assessment	11
1. Taxonomy	11
2. Pest Overview	12
3. Is the pest a vector?	16
4. Is a vector needed for pest entry or spread?	16
5. Regulatory status of the pest	16
6. Distribution	17
7. Habitats and their distribution in the PRA area	19
8. Pathways for entry	20
9. Likelihood of establishment in the natural environment (PRA area)	21
10. Likelihood of establishment in the managed environment (PRA area)	21
11. Spread in the PRA area	22
12. Impact in the current area of distribution	22
12.01. Impact on biodiversity and ecosystem patterns	22
12.02. Impact on ecosystem services	24
12.03. Socio-economic impact	24
13. Potential impact in the PRA area	25
14. Identification of the endangered area	26
15. Climate change	26
16. Overall assessment of risk	28
Stage 3: Pest risk management	30
17. Phytosanitary measures	30
17.01 Management measures for eradication, containment and control	31
18. Uncertainty	33
19. Remarks	34
20. References	35
Appendices	
Appendix 1 Projection of climate suitability for <i>Triadica sebifera</i>	40
Appendix 2 EU Biogeographical regions	53
Appendix 3 Images of <i>Triadica sebifera</i>	54
Appendix 4 Distribution summary for EU Member States and Biogeographical regions	57
Appendix 5 Distribution maps of <i>Triadica sebifera</i>	58

Summary¹ of the Express Pest Risk Analysis for *Triadica sebifera*

PRA area: The EPPO region (see https://www.eppo.int/ABOUT_EPPO/images/clickable_map.htm)

Describe the endangered area:

The Expert Working Group (EWG) considers that the endangered area is primarily coastal habitats, woodland and forests, grassland, land sparsely wooded and heathland within the Mediterranean and Black Sea biogeographic regions, including the following EU countries: Portugal, Spain, southern coast of France, Italy, Croatia, Greece. Turkey and Israel are also within the endangered area. The species distribution model (current climatic projection) predicts a region of potential suitability for *Triadica sebifera* in southern Europe (Appendix 1, Figure 5). Countries in which *T. sebifera* may be capable of establishing widely include all those bordering the northern Mediterranean Sea, from Portugal to Turkey, as well as the Black Sea coasts of Turkey, Georgia and Russia. The model predicts that establishment in the rest of Europe will largely be prevented because of low summer temperature, with moisture limitation in central Spain and frost limitation in far eastern central Europe (Appendix 1, Figure 6).

Within the EPPO region the species is currently absent from the natural environment. However, the EWG consider that the species will be able to establish within the EPPO region in similar habitats to that of its invasive range in the USA.

Main conclusions

The results of this PRA show that *Triadica sebifera* poses a high risk for biodiversity to the endangered area (Mediterranean and Black Sea biogeographical region) with a high uncertainty. *Triadica sebifera* is an aggressive invader in the southern United States and Australia. The most serious impacts of invasion appear to have occurred following widespread commercial and ornamental planting of the species (USDA 2008; Scheld and Cowles 1981; Bruce *et al.* 1997). While it displays broad habitat suitability, *T. sebifera* appears limited by cold winter temperatures and steep land gradients in the USA (Gan *et al.* 2009; USDA 2008). Warmer regions in the PRA area, specifically the Mediterranean, face the highest risk of *T. sebifera* establishment.

Entry

Plants for planting is the main pathway for entry into the EPPO region. The likelihood of entry is high with a moderate rating of uncertainty.

Establishment

The likelihood of establishment in the natural and managed environment is moderate with a high uncertainty for the former and moderate uncertainty for the latter. Temperature and precipitation are the most straightforward climatic factors to determine likelihood of establishment of the species in the non-native north American range. This is likely to be similar in the EPPO region. Areas that experience winters with temperatures regularly dipping below -12°C are most likely to constitute the northern limits of *T. sebifera* establishment in the EPPO region (Gan *et al.* 2009; Grace 2001). *Triadica sebifera* is not adapted to dry conditions and this can restrict the species establishment in dry habitats not close to waterbodies in the Mediterranean region.

Spread

Triadica sebifera has a high capacity of spread with low uncertainty. A variety of bird species (for example the European Starling) have been recorded consuming and spreading the seeds of *T. sebifera*, contributing to new establishments and the invasion success of *T. sebifera* in the Southern USA (Renne *et al.* 1999; 2001; 2002; Jubinsky and Anderson 1996). *Triadica sebifera* can also spread along waterways, after heavy rains, flooding, or proximity to rivers and streams as seeds can float (Jubinsky and Anderson 1996; Bruce *et al.* 1997; NSW Factsheet 2017). Deliberate planting of *T. sebifera* seeds or young plants remains the most likely form of human-assisted spread. *Triadica*

¹ The summary should be elaborated once the analysis is completed

sebifera has the potential to be spread through the horticultural trade where there are a number of suppliers selling the species within the EPPO region (for example <http://www.panglobalplants.com/>). The species is present in a limited number of botanical gardens within the EPPO region.

Potential impacts in the EPPO region

As *T. sebifera* is absent from the natural environment in the EPPO region, all data on impacts comes from the species' invasive range elsewhere. Thus, all information on impacts can only be used as a proxy to the EPPO region. The EWG consider impacts in the EPPO region will be similar to that seen in other invasive regions (USA and Australia). Impacts on biodiversity and ecosystem services will be high with a high level of uncertainty. Socio-economic impacts will be low with a moderate uncertainty. The habitats where the species has biodiversity and ecosystem service impacts in the USA are present in the EPPO region, including coastal habitats, woodland and forests (including riparian forests), grassland, land sparsely wooded and heathland. In similar climatic conditions to the EPPO region, such as California, the species has been recorded as naturalised with the potential to spread along low lying riparian habitats (Bower *et al.* 2009). On the basis of climatic and habitat similarity, similar impacts may be expected for the EPPO region.

In the most severely affected region, the Southern USA, it is possible that a high propagule pressure exacerbated the invasion. The USDA began promoting the large-scale planting of *T. sebifera* in the Gulf Coast states in the early 1900s. *Triadica sebifera* has been widely grown and planted for ornamental trade in Houston, Texas (USDA 2000; Carmarillo *et al.* 2015). Therefore, without this propagule pressure occurring in the EPPO region (the species is not considered a popular ornamental species), such impacts may not be realised.

In addition, warm, moist low-lying forest and coastal grasslands feature more prominently in the USA Gulf coast states than across the PRA region. The most severe impact of *T. sebifera* is the conversion of grassland to woody thickets.

Habitats where the species is most likely to establish within the EPPO region include: coastal habitats, woodland and forests, wet grassland, land sparsely wooded and heathland. *Triadica sebifera* will establish in a wide range of soil types: clays, loams, and sands.

The text within this section relates equally to EU Member States and non-EU Member States in the EPPO region.

Climate change

The climate change projections for the 2070s suggest that under the less extreme RCP4.5 scenario the suitable region in Europe will have extended northwards, especially in western Europe where warmer summers may mean that the southern United Kingdom and the southern Baltic coast become more suitable (Appendix 1, Figure 7). Under the more extreme RCP8.5 scenario, the northwards expansion is even greater, especially in eastern Europe (Appendix 1, Figure 8). This is likely driven by a relaxation of frost constraints.

In the evaluated climate change scenarios, predicted suitability was stable in the Mediterranean but increased in the Black Sea. Other biogeographic regions predicted to strongly increase in suitability are Atlantic, Continental, Pannonian and Steppic (Figure 9). The countries within the endangered area under climate change include: Portugal, Spain, France, Germany, Ukraine, Georgia, Turkey, Russia, Greece, Albania, Bosnia and Herzegovina, Croatia, Slovenia, Austria, Hungary, Italy, Israel and the north coastline of Algeria. The influence of projected climate change scenarios has not been considered in the overall scoring of the risk assessment based on the high levels of uncertainty with future projections.

The results of this PRA show that *Triadica sebifera* poses a high risk to the endangered area (Mediterranean and Black Sea biogeographical region) with a high uncertainty.

The major pathway(s) being considered:

(1) Plants for planting

Given the significant impact of the species in other parts of the world and the identified risk to the PRA area, the EWG recommends the following measures for the endangered area:

International measures:

For the pathway plants for planting:

- Prohibition of import into and movement within countries in the endangered area, of plants labeled or otherwise identified as *Triadica sebifera*,
- Recommend that *Triadica sebifera* is banned from sale within the endangered area,
- *Triadica sebifera* should be recommended for regulation within the endangered area.

National measures

Triadica sebifera should be monitored and eradicated, contained or controlled where it occurs in the endangered area. In addition, public awareness campaigns to prevent spread from existing populations or from botanic gardens in countries at high risk are necessary. If these measures are not implemented by all countries, they will not be effective since the species could spread from one country to another. National measures should be combined with international measures, and international coordination of management of the species between countries is recommended.

The EWG recommends the prohibition of selling and movement of the plant. These measures, in combination with management plans for early warning; obligation to report findings, eradication and containment plans, and public awareness campaigns should be implemented.

Containment and control of the species in the PRA area

Eradication measures should be promoted where feasible with a planned strategy to include surveillance, containment, treatment and follow-up measures to assess the success of such actions. As highlighted by EPPO (2009), regional cooperation is essential to promote phytosanitary measures and information exchange in identification and management methods. Eradication may only be feasible in the initial stages of infestation, and this should be a priority. The EWG considers that this is possible at the current level of occurrence the species has in the EPPO region.

General considerations should be taken into account for all potential pathways and these measures should involve awareness raising, monitoring, containment and eradication measures. NPPOs should facilitate collaboration with all sectors to enable early identification including education measures to promote citizen science and linking with universities, land managers and government departments.

Import for plant trade

Prohibition of the import, selling, planting, and movement of the plant in the endangered area.

Unintended release into the wild

The species should be placed on NPPO's alert lists and a ban from sale would be recommended in countries most prone to invasion. Export of the plant should be prohibited within the EPPO region.

Management measures would be recommended to include an integrated management plan to control existing populations including manual and mechanical techniques, targeted herbicides and proven biological control techniques. Monitoring and surveillance including early detection for countries most prone to risk. National Plant Protection Organisations (NPPOs) and other stakeholders should report any finding in the whole EPPO region in particular the Mediterranean and Black Sea.

Intentional release into the wild

Prohibition on planting the species or allowing the plant to grow in the wild.

Natural spread (method of spread within the EPPO region):

Increase surveillance in areas where there is a high risk the species may invade. NPPOs and other stakeholders should provide land managers and stakeholders with identification guides and facilitate regional cooperation, including information on site specific studies of the plant, control techniques and management.

See Standard PM3/67 ‘Guidelines for the management of invasive alien plants or potentially invasive alien plants which are intended for import or have been intentionally imported’ (EPPO, 2006).

<p>Phytosanitary risk (including impacts on biodiversity and ecosystem services) for the <u>endangered area</u> (current/future climate)</p> <p>Pathway for entry Plants for planting: High/Moderate Likelihood of establishment in natural areas: Moderate/High Likelihood of establishment in managed areas: Moderate/High Spread: High/High</p> <p>Impacts (EPPO region) Biodiversity: High/High Ecosystem services: High/High Socio-economic: Low/Moderate</p>	High X	Moderate	Low
<p>Level of uncertainty of assessment (current/future climate)</p> <p>Pathway for entry Plants for planting: Moderate/Moderate Likelihood of establishment in natural areas: High/High Likelihood of establishment in managed areas: Moderate/High Spread: Low/High</p> <p>Impacts (EPPO region) Biodiversity: High/High Ecosystem services: High/High Socio-economic: Moderate/High</p>	High X	Moderate	Low

Other recommendations:

The EWG recommend that *Triadica sebifera* is not utilised as a bioenergy crop within the EPPO region.

Express Pest Risk Analysis:

.....

(Triadica sebifera)

Prepared by: *S. Luke Flory and Austin Young, University of Florida, Gainesville, Florida, USA.*
flory@ufl.edu

Date: October 2017

Stage 1. Initiation

Reason for performing the PRA:

Triadica sebifera is currently absent from the natural environment in the European Union and the non-EU countries of the EPPO region. *Triadica sebifera* was included in a list of 95 invasive alien species that are likely to “arrive, establish, spread and have an impact on biodiversity or related ecosystem services in the EU over the next decade” (Roy *et al.* 2015). In 2016, the species was prioritized (along with 36 additional species from the EPPO List of Invasive Alien Plants and a recent horizon scanning study²) for PRA within the LIFE funded project “Mitigating the threat of invasive alien plants to the EU through pest risk analysis to support the Regulation 1143/2014” (see www.iap-risk.eu). *Triadica sebifera* was one of 16 species identified as having a high priority for PRA. In North America, *T. sebifera* readily invades a variety of natural habitats and disturbed sites, especially grasslands and areas adjacent to water (Gan *et al.* 2009, Pile *et al.* 2017). In these habitats, *T. sebifera* can dramatically alter community species composition, reduce diversity, and affect nutrient cycling (Pile *et al.* 2017). Species distribution modelling predicts that *T. sebifera* can establish in the EPPO region including EU Member States in northern Mediterranean biogeographical region, from Portugal to Turkey.

PRA area: The EPPO region (see https://www.eppo.int/ABOUT_EPPO/images/clickable_map.htm)

The risk assessments were prepared according to EPPO Standard PM5/5 (slightly adapted) which has been approved by the 51 EPPO Member Countries, and which sets out a scheme for risk analysis of pests, including invasive alien plants (which may be pests according to the definitions in the International Plant Protection Convention). EPPO engages in projects only when this is in the interests of all its member countries, and it was made clear at the start of the LIFE project that the PRA area would be the whole of the EPPO region. Furthermore, we believe that since invasive alien species do not respect political boundaries, the risks to the EU are considerably reduced if neighbouring countries of the EPPO region take equivalent action on the basis of broader assessments and recommendations from EPPO.

All information relating to EU Member States is included in the Pest risk analysis and information from the wider EPPO region only acts to strengthen the information in the PRA document. The PRA defines the endangered area where it lists all relevant countries within the endangered area, including EU Member States. The distribution section lists all relevant countries in the EPPO region (including by default those of EU Member States and biogeographical regions which are specific to EU member States). Habitats and where they occur in the PRA are defined by the EUNIS categorization which is relevant to EU Member States. Pathways are defined and relevant to the EU Member States and the wider EPPO Member countries, and where the EWG consider they may differ between EU Member States and non-EU EPPO countries, this is stated. The establishment and spread sections specifically detail EU Member States. When impacts are relevant for both EU Member States and non-EU EPPO countries this is stated ‘The text within this section relates equally to EU Member States and non-EU Member States in the EPPO region’.

²

<http://ec.europa.eu/environment/nature/invasivealien/docs/Prioritising%20prevention%20efforts%20through%20horizon%20scanning.pdf>

Where impacts are not considered equal to EU Member States and non-EU Member States this is stated and further information is included specifically for EU member States. For climate change, all countries (including EU Member States) are considered.

Stage 2. Pest risk assessment

1. Taxonomy:

Kingdom: Plantae, Tracheophyta, Spermatophytina, Magnoliopsida Rosanae, Malpighiales, Euphorbiaceae, Triadica Lour. Species: *Triadica sebifera* (L.) Small (ITIS Standard Report 2017).

Synonyms: *Croton sebiferum* L., *Sapium sebiferum* (L.) Roxb., *Stillingia sebifera* (L.) Michx., *Carumbium sebiferum* (L.) Kurz, *Excoecaria sebifera* (L.) Müll. Arg., *Seborium chinense* Raf.

This genus was for many years united with *Sapium* as sect. *Triadica*. Recently it has been considered to constitute a separate genus characterized by peculiar fruits, seeds, and leaf glands (Esser *et al.* 1997).

Common name:

English: Chinese tallow tree, Small – tallowtree, chicken tree; Florida aspen; popcorn tree; vegetable tallow; white wax berry

Spanish: arbol del sebo

French: arbre à suif; arbre savon; glutier

Chinese: wujiu

German: Chinesischer Talgbaum

Plant type: Deciduous medium sized tree species

Related species in the EPPO region: None

2. Pest overview

Introduction

Triadica sebifera is a deciduous tree up to 20 – 30 meters in height. It is native to China and Japan, but due to its commercial applications and popularity as an ornamental plant, *T. sebifera* has been introduced to Europe, North America, South America, Africa, Asia and the Australian continents (GBIF, EDDMAPS, CABI). After escape and spread, it is now recorded widely in the southern United States and Australia (NSW Department of Primary Industries, 2017; USDA 2008). Often, more than 95% of seeds are viable, but viability drops to between 10 and 50 % after one year in natural settings (Renne *et al.* 2002). A mature tree can produce an average of 100,000 seeds annually, depending on environmental conditions (USDA 2000; Jubinksy and Anderson 1996; Bruce *et al.* 1997). According to Scheld and Cowles (1981), *T. sebifera* can grow up to 2.8 m tall within two years after germination. In addition to intentional plantings by humans, birds and waterways disperse *T. sebifera* seeds (Bruce *et al.* 1997; Jubinksy and Anderson 1996; Renne *et al.* 1999). *Triadica sebifera* readily invades a variety of natural habitats and disturbed sites, especially grasslands and areas adjacent to water (Gan *et al.* 2009, Pile *et al.* 2017). In these habitats, *T. sebifera* can dramatically alter community species composition, reduce diversity, and affect nutrient cycling (Pile *et al.* 2017).

Environmental Requirements

Altitude

- In the southern forests of the United States approximately 80 % of tallow invasions occur at elevations lower than 50 m or slopes of $< 2^\circ$. At higher elevations and with steeper slopes, the likelihood of invasion drops dramatically and “no invasion was reported in the FIA [Forest Inventory Analysis] data for sites where elevation was greater than 165 m or slopes were steeper than 18° ” (Gan *et al.* 2009).
- In Taiwan, *T. sebifera* plantations occur at elevations of 400-700 m (Lin *et al.* 1958).
- Surveys of *T. sebifera* in the Indian Himalayas recorded trees at an altitude limit of 1,600 m (Jaryan *et al.* 2013).
- Pattison and Mack (2007) cites a record in the Himalayas of *T. sebifera* at 1,800 m, but the EWG was unable to retrieve the cited material.

Temperature

- In the southern forests of the United States *T. sebifera* does not occur at survey sites where the mean minimum temperature in January was below -12°C (10°F) (Gan *et al.* 2009).
- Grace (1998) suggested that the likeliest northern boundary of *T. sebifera* to be where average minimum winter temperatures dip to -12 to -15°C , or USDA zone 7b.
- In a climate modelling study, temperatures of 12°C and 24°C were determined as the lower temperature threshold for growth and lower limit of optimal temperature growth, respectively. The upper temperature limit for optimal growth and the upper temperature threshold for growth were determined to be 35°C and 40°C , respectively. They reached these values after consultation with average annual temperature data from the native range of *T. sebifera* (Pattison and Mack, 2008).

Precipitation

- In Taiwan, *T. sebifera* has been reported on sites with average annual precipitation ranging from 1,070 mm – 3,733 mm (Lin *et al.* 1958; Kirmse 1989; Meyer 2011).
- In the US and mainland China, 1,000 - 2,000 mm average annual precipitation (pers comm. E Siemann, 2017).
- Kirmse and Fisher (1989) described a New Mexico plantation (US) where *T. sebifera* grew well with average annual precipitation of 243 mm, although trees were irrigated for the first 2 months after transplanting (90.6% survival).

Soil

- Evidence from plantations and the native and naturalized range of *T. sebifera* indicates it is compatible with a wide range of soil types: clays, loams, and sands (Scheld and Cowles 1981; Bruce *et al.* 1995; Radford *et al.* 1968).
- *Triadica sebifera* is tolerant of saline soils (Scheld and Cowles 1981; Lin *et al.* 1958).
- In Taiwan, Lin *et al.* 1958 determined the chemical properties of soils on the “18 main habitats of *T. sebifera* tree.” Total nitrogen was 0-0.20% and pH was 2.9-8.5.
- *Triadica sebifera* growth, including height, diameter, aboveground biomass, shoot mass, and root mass was correlated with higher soil nutrition such as greater amounts of nitrogen, potassium, and phosphorus (Siemann and Rogers 2003; Rogers and Siemann 2002).
- Meyer (2011) summarized the findings of research on greater nitrogen content and the growth/establishment success of *T. sebifera*. The addition of nitrogen in soils enhances survival and growth but sometimes has no effect. It never has a negative effect.
- Gan *et al.* (2009) found that in sites throughout the Southeast USA, the probability of invasion was greater closer to water bodies, likely due in part to soil moisture levels that were favourable for establishment.
- *Triadica sebifera* exhibits significantly lower seedling survival in treatments with 23% soil moisture compared to treatments with $\geq 31\%$ soil moisture (Meyer 2011; Bruce *et al.* 1995).
- Pattison and Mack (2008) analysed moisture conditions in the native and naturalized ranges to determine optimum soil moisture conditions. They described these as 0.225 for the lower soil moisture survival threshold and 0.35 as the low soil moisture threshold for optimal growth (these values relate to the index of soil moisture used in CLIMEX when the soil moisture is to 0.2, this commonly results in grass wilting, whilst the value of 1 indicates the soil is saturated (Suthrest *et al.* 2004).

Light

- Jones and McLeod 1990 found a positive relationship between light availability and biomass production where the greatest production was achieved under 100 % light. In greenhouse experiments, seedling shoot and total biomass were significantly lower in ambient light than under an 87 % shade cloth. In the field, seedlings perform much better in open than shaded habitats.

Habitats

In North America, *T. sebifera* has a wide environmental tolerance and can thrive in many different habitats including forests (can invade closed and open forests), wetlands, grasslands, coastal prairie, mesic sites, disturbed sites, low-lying fields, swamp, and scrubby flatlands (Bruce *et al.* 1997; Camarillo *et al.* 2015; Langeland 2015). In the native range, in China, *T. sebifera* is found in disturbed habitats at low densities. In Australia, the species is reported from wet areas, along river, lakes, streams and swamps. However, it also grows in drier conditions including roadsides and disturbed areas (NWS Department of Primary Industries, 2017).

Symptoms (Impacts)

Where it invades, *T. sebifera* can disrupt ecosystem processes, decrease biodiversity, and alter community structure (Bruce *et al.* 1997; Jubinsky and Anderson 1996; Cameron and Spencer 1989). Due to its rapid leaf decomposition, *T. sebifera* can alter soil chemistry (Cameron and Spencer 1989), which may allow it to better compete with native plants (Camarillo *et al.* 2015). *Triadica sebifera* readily replaces native vegetation and can establish dominant stands (Bruce *et al.* 1997; Camarillo *et al.* 2015; Neyland and Meyer 1997). In Texas, invasions of *T. sebifera* have converted coastal grasslands into forests (Bruce *et al.* 1997; Camarillo *et al.* 2015). Harcombe *et al.* (1993) showed that 15-year-old *T. sebifera* stands had significantly higher net productivity than the ecosystem it replaced. Established stands of *T. sebifera* are very difficult to eradicate (USDA 2000; Camarillo *et al.* 2015; Bruce *et al.* 1997).

Identification

There are no congeneric species within the EPPO region including the EU Member States. A deciduous tree 20 to 30 meters tall (Appendix 3, Figure, 1). Low-spreading and multi-forked to tall and columnar. Stem often crooked and gnarled (Appendix 3, Figure, 2). Bark is grey, brown, and rough. Exudes a milky sap. Twigs slender. Petioles 2-6 cm long, with 2 sessile disc-shaped glands at the apex. Leaf-blades broadly rhombic-ovate, 2-7.5 x 1.5-7 cm, abruptly acutely acuminate, broadly cuneate to rounded, subtruncate at the base, entire, lateral nerves 7-12 pairs, glaucous beneath. Stipules 1-2 mm long, obtuse. *T. sebifera* flowers from April to June, producing both male and female flowers. Fruits are 1 cm three-lobed capsules expected to mature in the autumn. Seeds are 8 mm long and chalky white (covered by a white wax) (Flora of China 1972; Bruce *et al.* 1997; Scheld and Cowles 1981) (Appendix 3, Figure, 3). For additional information on identification including variations in morphology see Flora of China (1972).

Existing PRAs

Hawaii: Australia/New Zealand Weed Risk Assessment adapted for the Pacific. Pacific Island Ecosystems at Risk (PIER). This risk assessment predicts the likelihood of invasions of species in Australia, Hawaii, and the high islands of the Pacific. Results are also sometimes modified for the State of Florida. The risk assessment for Hawaii scored *T. sebifera* 14, indicating that the species poses a significant risk of becoming a problematic invader (PIER 2005). Accessed at: http://www.hear.org/pier/wra/pacific/triadica_sebifera_htmlwra.htm

Florida: Australia/New Zealand Weed Risk Assessment adapted for Florida. Data used for analysis published in: Gordon, D.R., D.A. Onderdonk, A.M. Fox, R.K. Stocker, and C. Gantz. 2008. Predicting Invasive Plants in Florida using the Australian Weed Risk Assessment. *Invasive Plant Science and Management* 1: 178-195. Score: 18 Accessed at: http://www.hear.org/wra/tncflwra/pdfs/tncflwra_sapium_sebiferum_ism.pdf

California: Bower M. J., Aslan C. E., Rejmánek M. 2009. Invasion potential of *T. sebifera* tree (*Triadica sebifera*) in California's Central Valley. *Invasive Plant Science and Management* 2: 386-395. This study determined California's Central valley to be highly susceptible to invasion by *T. sebifera*.

Georgia: The species was rejected with a high risk of invasiveness (score 19) and with a confidence of 75.8 out of 100. (https://pre.ice.ucdavis.edu/sites/default/files/pdf/farm_bill/PRE-5690.pdf)

Texas: The species was rejected with a high risk of invasiveness (score 20) and with a confidence of 81 out of 100.

Texas: https://pre.ice.ucdavis.edu/sites/default/files/pdf/farm_bill/PRE-5238.pdf

Spain: Andreu & Vilà (2009) performed two different types of Weed Risk Assessments (WRAs) for 80 species for Spain, including *T. sebifera*. For both the "Australian" WRA and "Weber-Gut" WRA methodologies *T. sebifera* earned an "intermediate" level of risk of invasion (Andreu & Vilà, 2009).

Socio-Economic Benefits

Triadica sebifera is a highly valued species for both its ornamental qualities and productive capability in agricultural and industrial sectors. Tallow wood is white and close-grained, suitable for carvings, furniture, carts, and match making as well as incense (Lin *et al.* 1958; USDA 2000). The leaves are used to make a black dye and manure (Lin *et al.* 1958; USDA 2000).

Triadica sebifera seeds are a source of vegetable tallow, a drying oil, and protein food. The outer covering of the seeds contains solid fat known as Chinese vegetable tallow and the kernels produce

a drying oil called Stillingia oil. Candles, soap, cloth dressing, and fuel are made from the tallow. The oil is used in machine oils, as a crude lamp oil, and in making varnishes and paints, because of its quick-drying properties. The presscake remaining after tallow and oil extraction can be processed to make a valuable animal feed and human food, rich in protein. Different parts of the plants are used in traditional Chinese medicine (USDA 2000, PROSEA 2001, Gao *et al.* 2016).

In the early 20th century, the Foreign Plant Introduction Division of the U.S. Department of Agriculture promoted tallow tree planting in the Gulf Coast and Southern US states to establish a local soap industry (Dewalt *et al.* 2011; USDA 2000). When cultivated under conventional agriculture methods, *T. sebifera* can be grown over extensive areas of land and can provide woody biomass for direct burning or conversion to charcoal, ethanol, and methanol (USDA 2000; Scheld and Cowles 1981). Wen *et al.* (2010) demonstrated *T. sebifera* seed oil can be converted to biodiesel. Due to its ability to re-sprout, its rapid growth rate, and its tolerance to flooding, drought and salt, *T. sebifera* has been considered as biomass crop for the Gulf coast region of the US (Scheld and Cowles 1981; USDA 2000).

Birds, wild and domestic, will feed on the seeds and it is sometime recommended as a bird food (USDA 2000, Renne *et al.* 1999). The flowers of *T. sebifera* are visited by honeybees, contributing to a commercially desirable, light-colored honey (Renne *et al.* 1999; USDA 2000). *Triadica sebifera* is recognized as an important species for commercial honey production in Louisiana (Lieux 1975).

Repeated introductions: *T. sebifera* has an extensive history of cultivation and sale and as an ornamental, especially in the in United States (Bruce *et al.* 1997, Camarillo *et al.* 2015, Siemann and Rogers 2003). By 1983, 200,000-300,000 trees were being grown for ornamental trade in Houston, Texas alone (USDA 2000).

Within the EPPO region, apart from the species in ornamental plant trade, there are no current socio-economic benefits for the species. Currently, there is little information available on the value of the species in horticulture. The species is listed as available from two suppliers in the UK (<https://www.rhs.org.uk/Plants/Nurseries-Search-Result?query=239677>). The EWG did not find any additional information for suppliers within the EPPO region including EU Member States except for the information below.

Seeds of the species are available via a number of online supplier:

<http://b-and-t-world-seeds.com/cartall.asp?species=Triadica%20sebifera&sref=545154>

<https://www.amazon.com/Chinese-Tallow-Triadica-sebifera-Ornamental/dp/B01L3A1DR4>

<http://www.panglobalplants.com/>

Historically at the beginning of the 19th century in Europe, many countries (for example Italy and France) have explored the economic benefits of the species for oil production (Antonelli, 1820; Société nationale d'horticulture de France., 1832; Loiseleur-Deslongchamps, 1832).

3. Is the pest a vector?

Yes No

In North America, *T. sebifera* is repeatedly described as relatively free of serious pests and even pest resistant (Rogers and Siemann 2002; Scheld and Cowles 1981; Jubinsky and Anderson 1996). No descriptions of negative ecological effects resulting from the transmission of a pest or pathogen into a novel environment by *T. sebifera* were found in the literature we reviewed.

4. Is a vector needed for pest entry or spread?

Yes No

No vector is necessary for *T. sebifera* to enter into or spread within the PRA area. Ornamental plantings, bird dispersal, and waterways, are primary mechanisms of seed dispersal (Bruce *et al.* 1997; Jubinsky and Anderson 1996).

5. Regulatory status of the pest

In the USA, *T. sebifera* is listed as a noxious weed in the following States: Florida, Mississippi, Texas, and Louisiana (<https://plants.usda.gov/core/profile?symbol=TRSE6>)

In Australia, *T. sebifera* is regarded as an environmental weed in New South Wales and a potential environmental weed in south-eastern Queensland (Brisbane City Council, 2018).

In the EPPO region there are no regulations specific to *T. sebifera*.

6. Distribution³

Continent	Distribution (list countries, or provide a general indication , e.g. present in West Africa)	Provide comments on the pest status in the different countries where it occurs (e.g. widespread, native, introduced....)	Reference
Africa	South Africa, Sudan, Uganda, Zambia	Non-native. Limited distribution. At this time, the EWG consider the species shows no signs of invasiveness in Africa.	CABI (2017), GBIF (2017), EDDMAPPS, USDA (2008);
Americas	USA (Mississippi, Texas, Florida, Georgia, South Carolina, North Carolina, California, Louisiana, Arkansas, Alabama, Hawaii, Louisiana, Oklahoma, Virginia) Mexico Caribbean (Martinique, Cuba) Brazil Peru	Non-native and invasive. Non-native and planted without showing signs of invasiveness	CABI (2017), GBIF (2017), EDDMAPPS, USDA (2008); Bruce (1997); Carmarillo <i>et al.</i> (2015), Siemann and Rogers (2003); Chapman <i>et al.</i> (2008); DeWalt <i>et al.</i> (2011)
Asia	China, Japan, South Korea India, Pakistan, Taiwan	Native Non-native and mostly planted. Taiwan (uncertain status)	CABI (2017), GBIF (2017), EDDMAPPS, USDA (2008), Bruce <i>et al.</i> (1997); Siemann and Rogers 2003;); DeWalt <i>et al.</i> (2011)
Europe	UK, France, Germany, the Netherlands, Italy, Portugal, Spain	Long history of planting within the region for ornamental purposes and other uses. Present in botanical gardens. Not present in the natural environment.	Antonelli, (1830), Banfi and Visconti (2013); Batchelor (2017); Dillwyn and Collinson (1843); Hortus Collinsonianus (1843); Hortus Paddingtonensis (1797); Hortus Kewensis (1814).
Oceania	Australia, New Zealand	Introduced and invasive in Australia New Zealand – planted individuals,	CABI (2017), GBIF (2017), EDDMAPPS, NSW Primary Industries (2017); Wang and Ecroyd (2009).

³ See also appendix 4: Distribution summary for EU Member States and Biogeographical regions

Continent	Distribution (list countries, or provide a general indication , e.g. present in West Africa)	Provide comments on the pest status in the different countries where it occurs (e.g. widespread, native, introduced....)	Reference
		not showing signs of invasiveness	

Introduction

Triadica sebifera is native to China and Japan with a non-native distribution showing invasive behaviour in the USA and Australia.

North America

In the Southern USA, *T. sebifera* is a major invasive species. Although it was first introduced in the late 18th century, it did not become invasive until the 20th century, especially the latter half. This invasion pattern may be due to greater propagule pressure brought about by large-scale commercial plantings and hurricanes. Invasions convert grasslands to woody thickets, displaces native species, and disrupts ecological processes. According to DeWalt *et al.* (2011): “Present populations near the sites of the earliest introductions of Chinese tallow tree to the southeastern USA (Charleston, South Carolina and Savannah, Georgia) appear to differ in genetic composition as well as genetic diversity from populations resulting from introductions made approximately 120 years later to the rest of the southeastern USA.”

Central and South America

In Central and South America, there is no evidence of widespread establishment or invasion by *T. sebifera*.

Australia

According to the NSW Government in Australia: “It is fast becoming an invasive environmental weed of water courses and native vegetation areas... Originally introduced to Australia as an ornamental tree with beautiful coloured foliage. It has been planted in streets and garden in southeast Queensland and northern New South Wales (NSW). Naturalised populations have now been identified in various locations throughout southeast Queensland. The largest infestation of *T. sebifera* tree exists near Casino, NSW. Smaller infestations are evident throughout the North Coast, Central Coast and New England regions of NSW. Localized plants also exist in Victoria.” (NSW Factsheet 2017). According to DeWalt *et al.* (2011): “We are unaware of when Chinese tallow was introduced to Australia; however, its genetic similarity to the rest of the southeastern USA indicates that it may have been introduced in the early 1900s to Australia at the same time it was being extensively planted in areas along the USA coast of the Gulf of Mexico or that it represents a secondary invasion from the southeastern USA.”

Europe

Triadica sebifera was introduced and planted in the south of France since the 19th century (Société nationale d'horticulture de France, 1832). The “Colonial Garden” of Paris tested the species for oil production during the beginning of 20th century. In the UK, it was introduced in the 18th century, probably as early as 1703 (Batchelor, 2017; Dillwyn and Collinson, 1843). In Italy, it was introduced and planted in Botanic Gardens and Agricultural experimental fields at least since the 19th century (Banfi and Visconti, 2013). In the Netherlands, it was introduced and planted in the

Botanic Garden of Amsterdam (pers. comm. G. Brundu, 2017). In Germany, Portugal and Spain, it was introduced and planted in Botanic Gardens (pers. comm. G. Brundu, 2017).

7. Habitats and their distribution in the PRA area

Habitats	EUNIS habitat types	Status of habitat (eg threatened or protected)	Is the pest present in the habitat in the PRA area (Yes/No)	Comments (e.g. major/minor habitats in the PRA area)	Reference
Coastal habitats	For example, B1.8 Moist and wet dune slacks	Partially threatened	No	Major	Pile <i>et al.</i> 2017
Mires, bogs, fens	For example, D5 Sedge and reed beds normally without free-standing water	Partially threatened	No	Medium	Pile <i>et al.</i> 2017
Woodland, forest and other wooded land	For example: G1, 1.1, 2, 3, 4 and 5	Partially threatened	No	Major	Pile <i>et al.</i> 2017
Grasslands	For example: E2, E3, E5 and E7	Partially threatened	No	Major	Pile <i>et al.</i> 2017
Habitat complexes	For example, X13 – 16, 24, 25	NA	No	Major	Pile <i>et al.</i> 2017
Heathland, scrub and tundra	For example: F9, FA & B	Partially threatened	No	Major	Pile <i>et al.</i> 2017

In North America, *T. sebifera* has a wide environmental tolerance and can thrive in many different habitats including forests (capable of invading closed and open forest systems), wetlands, grasslands, coastal prairie, mesic sites, disturbed sites, low-lying fields, swamp, and scrubby flatlands (Bruce *et al.* 1997; Camarillo *et al.* 2015; Langeland 2009). In the native range, in China, *T. sebifera* is found in disturbed habitats at low densities.

Within the EPPO region and Europe in particular, the species is currently absent from the natural environment. However, the EWG consider that the species will be able to establish within the EPPO region in similar habitats to that of its invasive range in the USA. These include coastal habitats, woodland and forests, grassland, land sparsely wooded and heathland. Other habitats detailed above can be at risk (see above).

8. Pathways for entry (in order of importance)

Possible pathways	Pathway: Plants and seeds for planting (CBD terminology: Escape from confinement)
Short description explaining why it is considered as a pathway	<p><i>T. sebifera</i> has a long history of deliberate planting within the EPPO region (see for example Kew, 1814; Antonelli, 1830). Seeds can be bought on the internet and shipped nearly anywhere in the EPPO zone (see below). <i>T. sebifera</i> is available via nurseries within the EPPO region. The species can also be purchased via online suppliers outside of the EPPO region.</p> <p>Examples of online suppliers include: http://b-and-t-world-seeds.com/cartall.asp?species=Triadica%20sebifera&sref=545154 https://www.amazon.com/Chinese-Tallow-Triadica-sebifera-Ornamental/dp/B01L3A1DR4 http://www.panglobalplants.com/</p>
Is the pathway prohibited in the PRA area?	No, the pathway is not prohibited within the EPPO region.
Has the pest already intercepted on the pathway?	Yes, the species is actively sold via a number of internet suppliers within the region and outside of the EPPO region (see above).
What is the most likely stage associated with the pathway?	All growth forms except for large trees are associated with this pathway, including trade of seeds.
What are the important factors for association with the pathway?	Nurseries and seed suppliers will ship stock throughout the world. The seed will survive storage.
Is the pest likely to survival transport and storage in this pathway?	Yes, seeds will survive storage and live plants will be shipped to ensure plant survival.
Can the pest transfer from this pathway to a suitable habitat?	Yes. Likely through intentional introductions by humans. However, given the seeds can be dispersed by birds and waterways, any tree producing berries within the EPPO region could be the origin of a new establishment, especially trees that are planted near moving bodies of water. <i>T. sebifera</i> can survive in urban, rural, natural, and disturbed habitat.
Will the volume of movement along the pathway support entry?	It is likely that all seed stock is obtained from outside of the EPPO region. However, the EWG does not have any figures on the volume of movement along this pathway.
Will the frequency of movement along the pathway support entry?	Yes, as it is likely that all seed stock is obtained from outside of the EPPO region. However, the EWG does not have any figures on the frequency of movement along this pathway.
Likelihood of entry	Low <input type="checkbox"/> Moderate High X
Likelihood of uncertainty	Low <input type="checkbox"/> Moderate X High <input type="checkbox"/>

As the species is imported as a commodity, all European biogeographical regions will have the same likelihood of entry and uncertainty scores.

Do other pathways need to be considered? NO

9. Likelihood of establishment in the natural environment PRA area

Temperature and precipitation are the most straightforward climatic factors to determine likelihood of establishment of the species in the non-native north American range. This is likely to be similar in the EPPO region including the EU territory. Areas that experience winters with temperatures regularly dipping below -12°C are most likely to constitute the northern limits of *T. sebifera* establishment in the EPPO region (Gan *et al.* 2009; Grace 1998). *Triadica sebifera* is not adapted to dry conditions and this can restrict the species establishment in habitats not close to waterbodies in the Mediterranean region.

Habitats where the species is most likely to establish within the EPPO region include: coastal habitats, woodland and forests, grassland, land sparsely wooded and heathland. *T. sebifera* will establish in a wide range of soil types: clays, loams, and sands.

The model predicts a region of potential suitability for *T. sebifera* in southern Europe (Appendix 1, Figure 5). Countries in which *T. sebifera* may be capable of establishing widely include all those bordering the northern Mediterranean Sea, from Portugal to Turkey. The model predicts that establishment in the rest of Europe will largely be prevented because of low summer temperature, with moisture limitation in central Spain and frost limitation in far eastern central Europe (Appendix 1, Figure 6).

In terms of Biogeographical Regions, those predicted to be most suitable for *T. sebifera* establishment in the current climate are the Mediterranean and Black Sea assuming the species is present close to waterbodies in micro-habitats with high soil moisture (Appendix 1, Figure 9).

Potential reasons why the species has not established within the EPPO region to-date, despite being planted historically may include, a lack of propagule pressure, lack of suitable genotypes, the species being contained when historically planted, and low precipitation or low temperatures specifically affecting seedling establishment.

There are no known natural enemies for this species in the EPPO region.

A moderate rating of likelihood of establishment has been given for the natural environment based on the lack of propagule pressure. A high rating of uncertainty has been given as the species is currently not present in the natural environment within the PRA area.

<i>Rating of the likelihood of establishment natural environment</i>	Low <input type="checkbox"/>	Moderate X	High <input type="checkbox"/>
<i>Rating of uncertainty</i>	Low <input type="checkbox"/>	Moderate <input type="checkbox"/>	High X

10. Likelihood of establishment in managed environment in the PRA area

Although *T. sebifera* is a popular ornamental species, there is no evidence that professional landscapers in Europe and the wider EPPO region plant this tree. The species has been grown within greenhouses in botanical gardens. *T. sebifera* is known to establish in disturbed environments in its invasive range (Renne *et al.* 1999; Jubinsky and Anderson 1996), so it is expected to be the same in the PRA area.

A moderate rating of likelihood of establishment in the PRA area in the managed environment has been given as the species has been shown to establish in these situations in similar climatic conditions to the EPPO region (EWG opinion). However, as this has yet to be realised in the EPPO region including the EU, a moderate rating of uncertainty is given.

<i>Rating of the likelihood of establishment in managed environment</i>	Low <input type="checkbox"/>	Moderate X	High <input type="checkbox"/>
<i>Rating of uncertainty</i>	Low <input type="checkbox"/>	Moderate X	High <input type="checkbox"/>

11. Spread in the PRA area

Natural Spread

A variety of bird species (for example the European Starling outside of the EPPO region) have been recorded consuming and spreading the seeds of *T. sebifera*, contributing to new establishments and invasion success of *T. sebifera* in the Southern US (Renne *et al.* 1999; 2001; 2002; Jubinsky and Anderson 1996). *Triadica sebifera* can also spread along waterways, after heavy rains, flooding, or proximity to rivers and streams because seeds can float (Jubinsky and Anderson 1996; Bruce *et al.* 1997; NSW Factsheet 2017). In the US, under natural means, *T. sebifera* has been shown to spread up to 350 km over a 20-year period (pers. comm. Siemann, 2017). Therefore, natural spread is likely to facilitate transfer to suitable habitats in the EPPO region including EU Member States. At present however, the volume of movement will not support spread within the PRA area as the species is not present in the natural environment.

Human assisted spread

Deliberate planting of *T. sebifera* seeds or young plants remains the most likely form of human assisted spread. *Triadica sebifera* has the potential to be spread through the horticultural trade as numerous suppliers selling the species within the EPPO region (for example <http://www.panglobalplants.com/>). The species is currently present in a limited number of botanical gardens within the EPPO region including EU Member States. Human assisted spread and the likelihood of transfer to a suitable habitat is high within the PRA area.

A high rating of spread has been given as the species has the potential to be spread by birds over long distances and via waterbodies, coupled with the movement of the species by humans.

<i>Rating of the magnitude of spread</i>	Low <input type="checkbox"/>	Moderate <input type="checkbox"/>	High X
<i>Rating of uncertainty</i>	Low X	Moderate <input type="checkbox"/>	High <input type="checkbox"/>

12. Impact in the current area of distribution

12.01 Impacts on biodiversity

In the USA, *T. sebifera* displaces native plant species and establishes dominant stands, and can transform areas of prairie and grassland to woody thickets within ten years (Bruce *et al.* 1995, Bruce *et al.* 1997, Siemann and Rogers 2003, Cameron and Spencer 1989). *Triadica sebifera* has been shown to invade south east coastal prairie that is the sole habitat of the federally endangered Attwater's prairie chicken (*Tympanuchus cupido attwateri*), the exclusive wintering ground of the federally endangered whooping crane (*Grus americana*), and important habitat for several other critically imperilled grassland birds. In addition, one federally endangered and 12 critically imperilled (category 11) plant species occur in the remaining fragments of this once vast system.

When leaf material becomes incorporated into the water body, this has shown to be toxic to amphibians (Cotton *et al.* 2012). Leonard (2008) showed that the leaf litter can impact the native habitat by: (1) Leaf litter has a direct effect on water quality (2) *T. sebifera* can cause differential survival and performance of tadpoles (3) Differences in water quality due to leaf litter can cause

changes in tadpole behavior (4) *T. sebifera* leaf litter breaks down much faster than litter from native trees and this influences aquatic community composition.

Triadica sebifera exhibits rapid leaf decay, which may alter soil nutrients (primarily higher levels of nitrogen) in ecosystems with fewer deciduous trees. *Triadica sebifera* can alter nutrient cycling and the species composition of decomposers.

Triadica sebifera supports a lower diversity of arthropods than co-occurring native trees in Texas (Hartley *et al.* 2010) and native prairies it displaces (Hartley *et al.* 2004, Cameron and Spencer 2010).

The NSW Department of Primary Industries (2017) highlights that impacts seen in the US are similar to that seen in Australia.

At 15 years old, *T. sebifera* stands in Texas demonstrated significantly higher net primary productivity than adjacent native prairie (Harcombe *et al.* 1993). Cameron and Spencer (1989) found that in *T. sebifera* woodland the concentration of nutrients P, K, NO₃, N, Zn, Mn, and Fe were significantly higher than native prairie, while Mg and Na were significantly lower. *T. sebifera* exhibits rapid leaf decay, which may increase nutrient alteration in ecosystems with fewer deciduous trees (Cameron and Spencer 1989).

In invasive contexts, *T. sebifera* can alter nutrient cycling and species composition among decomposers (Weber 2003; Zou *et al.* 2006; Cameron and Spencer 1989). Siemann and Rogers (2003) demonstrated that the competitive advantage displayed by *T. sebifera* in the Southern US may be due to escape from specialist herbivores. After comparing populations from the invasive range in Texas to populations from the native range in China, Siemann and Rogers (2005) concluded that “invasive *Sapium* ecotypes have a greater capacity to compensate for herbivory damage than native *Sapium* ecotypes.”

Based on the impacts shown in the current area of distribution, a high rating of impact on biodiversity has been given with low uncertainty.

<i>Rating of the magnitude of impact in the current area of distribution</i>	Low <input type="checkbox"/>	Moderate <input type="checkbox"/>	High X
<i>Rating of uncertainty</i>	Low X	Moderate <input type="checkbox"/>	High <input type="checkbox"/>

12.01. Impacts on ecosystem services

Ecosystem service	Does the IAS impact on this Ecosystem service? Yes/No	Short description of impact	Reference
Provisioning	Yes	<i>T. sebifera</i> can decrease water quality. The species can also displace native species reducing genetic resources.	Leonard 2008
Regulating	Yes	<i>T. sebifera</i> seedlings can establish at such high densities that fine fuels are lacking even in a stand of smaller trees. These changes in fuel characteristics result in patchier and/or less severe fires, negatively impacting native species (Meyer 2011). <i>T. sebifera</i> may render some areas inflammable. <i>T. sebifera</i> stands in Texas demonstrated higher net productivity than the original grassland ecosystem.	Meyer 2011; Grace 1998, Weber 2003; Zou <i>et al</i> 2006; Cameron and Spencer 1989; Harcombe <i>et al.</i> 1993
Cultural	Yes	Transforms grassland into woody thickets which can reduce human access to areas for recreational purposes and possibly adjacent waterways.	Bruce <i>et al.</i> 1997; Jubinsky and Anderson 1996

Based on the impacts shown in the current area of distribution, a high rating of impact has been given with low uncertainty.

Rating of the magnitude of impact in the current area of distribution	Low <input type="checkbox"/>	Moderate <input type="checkbox"/>	High <input checked="" type="checkbox"/>
Rating of uncertainty	Low <input checked="" type="checkbox"/>	Moderate <input type="checkbox"/>	High <input type="checkbox"/>

12.02. Describe the adverse socio-economic impact of the species in the current area of distribution

Removal of large trees is expensive (mulching can cost up to \$1 000 per acre) and *T. sebifera* will regenerate if not fully excavated (Jubinsky and Anderson, 1996, USDA, 2000, Meyer, 2011). However, current management practices are to apply herbicide to cut stumps. In addition, following up monitoring will be needed to exhaust the seed bank in areas under control. This will incur additional costs. In the USA, the species has been shown to degrade pasture land (Pile *et al.* 2017).

Human contact with the sap can causes irritation and ingesting any part of this plant can cause gastrointestinal upset, nausea, and vomiting (USDA, 2000). The species is also toxic to livestock (Russell *et al.* 1969).

Based on the cost of removal and the fact that the species has been shown to cause impacts to human and livestock a moderate rating of socio-economic impact has been given, with low uncertainty.

<i>Rating of the magnitude of impact in the current area of distribution</i>	Low <input type="checkbox"/>	Moderate X	High <input type="checkbox"/>
<i>Rating of uncertainty</i>	Low X	Moderate <input type="checkbox"/>	High <input type="checkbox"/>

13. Potential impact in the PRA area

As *T. sebifera* is absent from the natural environment in the EPPO region, all data on impacts comes from other regions of the invaded range. Thus, all information on impacts can only be used as a proxy to the EPPO region including the EU territory.

Will impacts be largely the same as in the current area of distribution? **Yes (in part)**

The habitats where the species has biodiversity and ecosystem service impacts in the US are present in the EPPO region, and EU territory in particular, including coastal habitats, woodland and forests (including riparian forests), grassland, land sparsely wooded and heathland. In similar climatic conditions to the PRA area, such as California, the species has been recorded as naturalised with the potential to spread along low lying riparian habitats (Bower *et al.* 2009). On the basis of climatic and habitat similarity, similar impacts may be expected for the EPPO region and the and EU territory.

However, in the most severely affected region, the Southern USA, it is possible that a high propagule pressure exacerbated the impact. The USDA began promoting the large-scale planting of *T. sebifera* in the Gulf Coast states in the early 1900s (though potentially as early as the 1700s (Pile *et al.* 2017). *T. Triadica sebifera* has been widely grown and planted for ornamental trade in Houston, Texas alone (USDA 2000; Carmarillo *et al.* 2015). Therefore, without this propagule pressure occurring in the EPPO region, such impacts may not be realised.

In addition, warm, moist low-lying forest and coastal grasslands feature more prominently in the USA Gulf States than across the PRA region. The most severe impact of *T. sebifera* is the conversion of grassland to woody thickets.

Based on climate modelling (see appendix 1), the areas that would face the greatest risk of severe impact are wet grasslands and riparian woodlands in the Mediterranean and Black Sea.

The text within this section relates equally to EU Member States and non-EU Member States in the EPPO region.

13.01. Negative impacts on biodiversity

A higher rating for uncertainty (compared to the current area of distribution) reflects the fact that the species is not currently present in the natural environment within the EPPO region and EU Member States.

<i>Rating of the magnitude of impact in the area of potential establishment</i>	Low <input type="checkbox"/>		Moderate <input type="checkbox"/>	High X
<i>Rating of uncertainty</i>	Low		Moderate <input type="checkbox"/>	High X

13.02. Negative impact the pest may have on categories of ecosystem services

A higher rating for uncertainty (compared to the current area of distribution) reflects the fact that the species is not currently present in the natural environment within the EPPO region and EU Member States.

<i>Rating of the magnitude of impact in the area of potential establishment</i>	<i>Low</i> <input type="checkbox"/>	<i>Moderate</i> <input type="checkbox"/>	<i>High</i> X
<i>Rating of uncertainty</i>	<i>Low</i>	<i>Moderate</i> <input type="checkbox"/>	<i>High</i> X

13.03 Socio-economic impact of the species

A lower rating of socio-economic impact and a higher rating of uncertainty (compared to the current area of distribution) reflects the fact that the species is not present in the EPPO region and EU Member States and thus management costs and other associated costs would not be as significant as in the invaded range.

<i>Rating of the magnitude of impact in the area of potential establishment</i>	<i>Low</i> X	<i>Moderate</i>	<i>High</i> <input type="checkbox"/>
<i>Rating of uncertainty</i>	<i>Low</i>	<i>Moderate</i> X	<i>High</i> <input type="checkbox"/>

14. Identification of the endangered area

The EWG considers that the endangered area is primarily woodland and forests, grassland, land sparsely wooded, heathland and dune slacks within the Mediterranean and Black Sea biogeographic regions. Within this area, habitats most susceptible are those with sufficient soil moisture such as margins of small waterbodies (e.g. rivers, ponds or lakes). Although there is limited suitability in other regions, e.g. the Atlantic areas, the EWG considers that these areas are less likely to be at risk from invasion.

The model predicts a region of potential suitability for *T. sebifera* in southern Europe (Figure 5). Countries in which *T. sebifera* may be capable of establishing widely include all those bordering the northern Mediterranean Sea, from Portugal to Turkey. The model predicts that establishment in most of the rest of Europe (see Figure 6) is less likely because of low summer temperature. In addition, moisture is expected to be limiting in central Spain and frost may reduce suitability in far eastern central Europe (Figure 6).

In terms of Biogeographical Regions those predicted to be most suitable for *T. sebifera* establishment in the current climate are the Mediterranean and Black Sea biogeographical regions, including the following countries EU: Portugal, Spain, southern coast of France, Italy, Croatia, and Greece.

15. Climate change

The influence of projected climate change scenarios has not been taken into account in the overall scoring of the risk assessment based on the high levels of uncertainty with future projections.

The climate change projections for the 2070s suggest that under the less extreme RCP4.5 scenario the suitable region in Europe will have extended northwards, especially in western Europe where warmer summers may mean that the southern United Kingdom and the southern Baltic coast become more suitable (Figure 7). Under the more extreme RCP8.5 scenario, the northwards expansion is even greater, especially in eastern Europe (Figure 8). This is likely driven by a relaxation of frost constraints.

In the evaluated climate change scenarios, predicted suitability was stable in the Mediterranean but increased in the Black Sea. Other biogeographic regions predicted to strongly increase in

suitability are Atlantic, Continental, Pannonian and Steppic (Figure 9). The climate change projections for the 2070s suggest that under the less extreme RCP4.5 scenario the suitable region in Europe will have extended northwards, especially in western Europe where warmer summers mean the species may be capable of establishing in southern United Kingdom and the southern Baltic coast (Figure 7). Under the more extreme RCP8.5 scenario, the northwards expansion is even greater, especially in eastern Europe (Figure 8). This is likely driven by a relaxation of frost constraints. The countries within the endangered area under climate change include: Portugal, France, Germany, Ukraine, Georgia, Turkey, Russia, Greece, Albania, Bosnia and Herzegovina, Croatia, Slovenia, Austria, Hungary, Italy, and the north coastline of Algeria.

15.01. Define which climate projection you are using from 2050 to 2100

Climate projection: 2070

15.02 Which component of climate change do you think is most relevant for this organism?

Temperature (yes)	Precipitation (yes)	CO ₂ levels (no)
Sea level rise (no)	Salinity (no)	Nitrogen deposition (yes)
Acidification (no)	Land use change (yes)	Other (please specify)

Are the introduction pathways likely to change due to climate change? (If yes, provide a new risk and uncertainty score)	Reference
No. The introduction pathways are unlikely to change (moderate with moderate uncertainty) as a result of climatic change because the primary pathway for entry into the EPPO region is intentional landscape plantings.	EWG opinion
Is the risk of establishment likely to change due to climate change? (If yes, provide a new risk and uncertainty score)	Reference
Yes. The area climatically suitable for <i>T. sebifera</i> will increase, therefore the risk of establishment in the overall EPPO region will increase as well. If precipitation increases due to climate change <i>T. sebifera</i> could increase the possibility of establishment. The rating for establishment is increased from moderate to high with high uncertainty for the natural environment and from moderate to high with a high uncertainty for the managed environment.	Wang <i>et al.</i> 2011; Gan <i>et al.</i> 2009
Is the risk of spread likely to change due to climate change? (If yes, provide a new risk and uncertainty score)	Reference
In the PRA region, winter temperature is a major limiting factor for the establishment and spread of <i>T. sebifera</i> . If average winter temperatures increase due to climate change, potential range of <i>T. sebifera</i> could increase as well. The rating of spread will remain high and the uncertainty will raise from low to high	(See climate modelling appendix)
Will impacts change due to climate change? (If yes, provide a new risk and uncertainty score)	Reference
With increasing temperature it is possible <i>T. sebifera</i> may impact the EPPO region as the area of suitability increases. Given the majority of suitable land in the EPPO region is at the northern limits of <i>T. sebifera</i> establishment, higher temperatures could cause more	EWG opinion

rapid growth and biomass accumulation, resulting in greater impacts to native species.

However, the rating for impact on biodiversity and ecosystem services will remain high with high uncertainty and socio-economic impacts will increase from low to moderate with a high uncertainty.

16. Overall assessment of risk

The results of this PRA show that *Triadica sebifera* poses a high risk to biodiversity in the endangered area (Mediterranean and Black Sea biogeographical region) with a high uncertainty. *Triadica sebifera* is an aggressive invader in the southern United States. The most serious effects of invasion appear to have occurred following widespread commercial and ornamental planting of the species (USDA 2008; Scheld and Cowles 1981; Bruce *et al.* 1997). While it displays broad habitat suitability, *T. sebifera* appears limited by cold winter temperatures and steep gradients (Gan *et al.* 2009; USDA 2008). Warmer regions in the PRA area, specifically the Mediterranean, face the highest risk of *T. sebifera* establishment. However, there is currently no evidence of *T. sebifera* establishment in the PRA region.

Pathways for entry:

Plants for planting

Likelihood of entry	Low	Moderate	High X
Likelihood of uncertainty	Low	Moderate X	High

Likelihood of establishment in the natural environment in the PRA area

Rating of the likelihood of establishment in the natural environment	Low	Moderate X	High
Rating of uncertainty	Low	Moderate	High X

Likelihood of establishment in managed environment in the PRA area

Rating of the likelihood of establishment in the managed environment	Low	Moderate X	High
Rating of uncertainty	Low	Moderate X	High

Spread in the PRA area

Rating of the magnitude of spread	Low	Moderate	High X
Rating of uncertainty	Low X	Moderate	High

Impacts

Impacts on biodiversity and the environment

Rating of the magnitude of impact in the current area of distribution	Low	Moderate	High X
Rating of uncertainty	Low X	Moderate	High

Impacts on ecosystem services

Rating of the magnitude of impact in the current area of distribution	Low	Moderate	High X
Rating of uncertainty	Low X	Moderate	High

Socio-economic impacts

Rating of the magnitude of impact in the current area of distribution	Low	Moderate X	High
Rating of uncertainty	Low X	Moderate	High

Impacts in the PRA area

Will impacts be largely the same as in the current area of distribution? **Yes (in part)**

Impacts on biodiversity in the PRA area

<i>Rating of magnitude of impact on biodiversity in the area of potential establishment</i>	<i>Low</i> <input type="checkbox"/>	<i>Moderate</i> <input type="checkbox"/>	<i>High</i> X
<i>Rating of uncertainty</i>	<i>Low</i>	<i>Moderate</i> <input type="checkbox"/>	<i>High</i> X

Impact on ecosystem services in the PRA area

<i>Rating of magnitude of impact on ecosystem services in the area of potential establishment</i>	<i>Low</i> <input type="checkbox"/>	<i>Moderate</i> <input type="checkbox"/>	<i>High</i> X
<i>Rating of uncertainty</i>	<i>Low</i>	<i>Moderate</i> <input type="checkbox"/>	<i>High</i> X

Socio-economic impact in the PRA area

<i>Rating of magnitude of socio-economic impact in the area of potential establishment</i>	<i>Low</i> X	<i>Moderate</i> <input type="checkbox"/>	<i>High</i>
<i>Rating of uncertainty</i>	<i>Low</i>	<i>Moderate</i> X	<i>High</i>

Stage 3. Pest risk management

17. Phytosanitary measures to prevent entry

The results of this PRA show that *Triadica sebifera* poses a high risk to the endangered area (Mediterranean and Black Sea biogeographical region) with a high uncertainty.

The major pathway(s) being considered:

(1) Plants for planting

Given the significant impact of the species in other parts of the world and the identified risk to the PRA area, the EWG recommends the following measures for the endangered area:

International measures:

For the pathway plant for planting:

- Prohibition of import into and movement within countries in the endangered area, of plants labeled or otherwise identified as *T. sebifera*,
- Recommend that *T. sebifera* is banned from sale within the endangered area,
- *Triadica sebifera* should be recommended for regulation within the endangered area.

National measures

Triadica sebifera should be monitored and eradicated, contained or controlled where it occurs in the endangered area. If these measures are not implemented by all countries, they will not be effective since the species could spread from one country to another. National measures should be combined with international measures, and international coordination of management of the species between countries is recommended.

The EWG recommends the prohibition of selling and movement of the plant. These measures, in combination with management plans for early warning; obligation to report findings, eradication and containment plans, and public awareness campaigns should be implemented.

Containment and control of the species in the PRA area

Eradication measures should be promoted where feasible with a planned strategy to include surveillance, containment, treatment and follow-up measures to assess the success of such actions. As highlighted by EPPO (2009), regional cooperation is essential to promote phytosanitary measures and information exchange in identification and management methods. Eradication may only be feasible in the initial stages of infestation, and this should be a priority. The EWG considers that this is possible at the current level of occurrence the species has in the EPPO region.

General considerations should be taken into account for all potential pathways and these measures should involve awareness raising, monitoring, containment and eradication measures. NPPOs should facilitate collaboration with all sectors to enable early identification including education measures to promote citizen science and linking with universities, land managers and government departments.

Import for plant trade

Prohibition of the import, selling, planting, and movement of the plant in the endangered area.

Unintended release into the natural environment

The species should be placed on NPPO's alert lists and a ban from sale would be recommended in countries most prone to invasion. Export of the plant should be prohibited within the EPPO region. Management measures would be recommended to include an integrated management plan to control including manual and mechanical techniques, targeted herbicides and proven biological control techniques. Monitoring and surveillance including early detection for countries most prone to risk. NPPOs should report any finding in the whole EPPO region in particular the Mediterranean and Black Sea area.

Intentional release into the natural environment

Prohibition on planting the species or allowing the plant to grow in the wild.

Natural spread (method of spread within the EPPO region):

Increase surveillance in areas where there is a high risk the species may invade. NPPO's should provide land managers and stakeholders with identification guides and facilitate regional cooperation, including information on site specific studies of the plant, control techniques and management.

See Standard PM3/67 'Guidelines for the management of invasive alien plants or potentially invasive alien plants which are intended for import or have been intentionally imported' (EPPO, 2006).

17.01 Management measures for eradication, containment and control

Management measures for control (Source: FLEPPC Management Plan 2005, Source: NSW Factsheet 2017)

Mechanical: Most sources describe *T. sebifera* infestations as nearly impossible to eliminate. Mechanical control of *T. sebifera* in Florida have either failed or exacerbated the infestation (FLEPPC 2005). Previous attempts at control have been through application of prescribed burns and mechanical removal of trees and seedlings, but there is limited success due to the density of growth and likelihood of disturbing non-target vegetation and soil. If the roots are not completely removed, the tree will regenerate. However, most sources agree that control should take place over a 3-5 year period for best results. (Bruce *et al.* 1997; Jubinsky and Anderson 1996; Meyer 2011, Pile *et al.* 2017)

Biological: Proven, reliable, biological control agents are not presently available. However, herbivores may be an important part of controlling *T. sebifera*. Siemann and Rogers (2003a/c) compared growth and herbivory levels in tallow trees cultivated from native (Asian) and invasive (Texas and Hawaii) seed grown in competitive common gardens in Texas and Hawaii. According to a FLEPPC (2005) summary of their results:

"Their results show that in the Texas garden, Asian genotypes were significantly smaller than North American genotypes, and both genotypes suffered very low levels of insect damage. However, in the Hawaiian garden, the situation was reversed. North American genotypes were significantly smaller than Asian genotypes. Additionally, North American genotypes experienced significantly more insect damage than Asian genotypes in the Hawaiian garden. The herbivore responsible for most foliar damage was *Adoretus sinicus* Burmsiter (Coleoptera: Scarabaeidae), a generalist chewing herbivore known to feed on more than 250 plant species in Hawaii (Siemann and Rogers, 2003b). The study demonstrates that invasive species liberated from herbivory may evolve greater

competitive ability, allowing them to expand their ranges. However, in the presence of an abundant generalist herbivore from the native range, this did not occur.”

There are no known natural enemies for this species in the EPPO region.

This section details the chemical control options utilised throughout the introduced range. Any detail of a product does not imply the product is legal or safe to use in the EPPO region.

Chemical: In Florida, basal bark herbicide applications are commonly used for control. According to Jubinsky (1996), herbicide control can be achieved with GARLON 4 and a spray adjuvant formulated to mix with oil-soluble herbicide.

The Florida Exotic Pest Plant Council recommends unique chemical control strategies for home sites and large areas:

Specific recommendations for home sites: Homeowners with one or a few trees should use Brush-B-Gon™ or Brush Killer™ herbicide. These diluted herbicide products (8.0% and 8.8% triclopyr amine, respectively) are available in quart-size containers from retail nursery supply stores. Property owners with larger stands of trees can use the more concentrated Garlon 3A™ or Garlon 4™ (44.4% triclopyr amine and 61.6% triclopyr ester, respectively), available in 2.5-gallon or larger containers from farm supply stores. If the dead trees will remain standing, Garlon 4™ can be diluted at a rate of one part herbicide to five parts oil (i.e., 20%) and applied to the bark at the base of trees with stems less than 6 inches in diameter

Specific recommendations for large areas: Make basal bark application, according to label instructions, of triclopyr ester- containing herbicide such as Garlon 4™ or Pathfinder IITM or cut stump application of triclopyr ester or triclopyr amine containing herbicide such as Garlon 3A™. Use at least 20% dilution in oil for basal bark application and at least 10% dilution (water for triclopyr amine) for cut stump application. For cut stump application, apply herbicide solution immediately after cutting. Coppicing may be reduced by addition of imazapyr-containing herbicide (e.g., 3% Stalker™ or Arsenal™). Seedlings and saplings can be controlled using foliar application (July to October) of imazapyr-containing herbicide (1% water-solution of Arsenal™ or Habitat™), fosamine-containing herbicide (30% water solution of Krenite STM), or triclopyr ester-containing herbicide (2% water-solution of Garlon 4™). For trees growing in water, aquatic labeled herbicides such as Habitat™ (imazapyr) or Renovate™ (triclopyr ester) must be used.

Australian weed authorities suggest cut stump or stem injection applications with the following herbicides:

Glyphosate 360 g/L with Metsulfuron-methyl 600 g/kg (Various products)

Rate: Tank mix of 1:1.5 of glyphosate plus 1 g of metsulfuron-methyl in 1 L of water

Comments: Stem injection method.

Withholding period: Nil.

Herbicide group: M, Inhibitors of EPSP synthase

Resistance risk: Moderate

Picloram 44.7 g/kg + Aminopyralid 4.47 g/L (Vigilant II ®)

Rate: Undiluted

Comments: Cut stump/stem injection application. Apply a 3–5 mm layer of gel for stems less than 20 mm. Apply 5 mm layer on stems above 20 mm.

Withholding period: Nil.

Herbicide group: I, Disruptors of plant cell growth (synthetic auxins) Resistance risk: Moderate

It should be highlighted that the availability of products containing these active substances will vary nationally and other products may be available and effective. Indications of the approved uses for each active substance may be incomplete. Products should be used following the instructions on the label and in line with the relevant plant protection product regulations.

As there are no occurrences of *T. sebifera* in the EPPO region (and the EU) in the natural environment, implementation costs for Member States would be relatively low. The cost of inaction could significantly increase potential costs in the future as any management programme would have to take place on a larger scale and this would reduce the cost-effectiveness of any measures.

18. Uncertainty

There is a high uncertainty for establishment in the natural environment within the EPPO region. The species has historically been planted within the EPPO region but it has not shown any signs of invasiveness. In the US, establishment has been aided by high propagule pressure where the USDA began promoting the large-scale planting of *T. sebifera* in the Gulf Coast states in the early 1900s. *T. sebifera* was widely planted as an ornamental in Houston during the 1950-1980s (USDA 2000; Carmarillo *et al.* 2015). Therefore, without this propagule pressure occurring in the EPPO region, such impacts may not be realised. Finally, a high level of uncertainty has been highlighted for impacts on biodiversity and ecosystem services. This uncertainty is linked to the uncertainty in establishment and the low propagule pressure in the absence of widespread planting. The high uncertainty of invasion risk within the PRA area is further supported by a lack of information concerning the source and amount of plant material introduced from outside of the EPPO region.

Uncertainty relating to the modelling includes:

Modelling the potential distributions of range-expanding species is always difficult and uncertain. We did not have sufficient information to determine whether all records used in the modelling were from established populations rather than plantations. If the latter were included, management to alleviate climatic stresses (such as irrigation) may have caused the model to over-estimate the niche breadth and potential establishment distribution of *T. sebifera*.

Triadica sebifera is largely restricted to wet micro-habitats such as river banks in the more arid parts of its global distribution (e.g. California and Mexico), so it is likely that establishment in the more arid parts of the suitable region in Europe would also be restricted to such habitats. Although we attempted to model this interaction by including river density in the model, local habitat factors are unlikely to be well represented at the scale of the model.

The limiting factors map may have under-estimated the limiting influence of winter temperatures in Europe, since two of the algorithms in the ensemble did not model a strong limitation of suitability at very cold temperatures. This will have raised the ensemble model suitability response to very cold winter temperatures.

Other variables potentially affecting the distribution of the species, such as edaphic variables, were not included in the model.

To remove spatial recording biases, the selection of the background sample was weighted by the density of Tracheophyte records on the Global Biodiversity Information Facility (GBIF). While this is preferable to not accounting for recording bias at all, a number of factors mean this may not be the perfect null model for species occurrence:

- The GBIF API query used to did not appear to give completely accurate results. For example, in a small number of cases, GBIF indicated no Tracheophyte records in grid cells in which it also yielded records of the focal species.
- Additional data sources to GBIF were used, which may have been from regions without GBIF records.

19. Remarks

The EWG recommend that *Triadica sebifera* is not utilised as a bioenergy crop within the EPPO region. At the present time this has not be realised in the EPPO region. Low and Booth (2007) state: US Biofuel company AgriBioFuels claims that Chinese tallow can produce 500 gallons of oil per acre, compared to 48 gallons for soybeans. They are working with Texas A&M Research Centre to develop the plant as a crop. Oil company Chevron has built a biodiesel plant in Texas, and Chinese tallow is considered a major potential feedstock. There has also been interest in Australia. A 2001 report on biofuels by the Western Australian Department of Agriculture recommended Chinese tallow as one of several oil-bearing plants “that should undergo some preliminary studies in Western Australia”. Its potential has been discussed on the Biofuels Forum, the forum of the Australian Biofuel Users group.

20. References

- Andreu J, Vilá M (2009) Risk analysis of potential invasive plants in Spain. *Journal of Nature Conservation*, doi:10.1016/j.jnc.2009.02.002. Book digitized by Google. Source: <https://books.google.it/books?id=P31LAAAAcAAJ>
- Antonelli G (1830) Nuovo Dizionario universale tecnologico o di arti e mestieri e della economia industriale e commerciale compilato dai signori Lenormand etc. Prima trad. italiana, Volume 1.
- Banfi E, Visconti A (2013) L'Orto di Brera alla fine della dominazione asburgica e durante l'età napoleonica, *Atti Soc. it. Sci. nat. Museo civ. Stor. nat. Milano*, 154 (II): 173-264
- Batchelor R (2017) John Bradby Blake, the Chinese tallow tree and the infrastructure of botanical experimentation. *Curtis's Botanical Magazine* 2017 vol. 34 (4): pp. 402–426
- Bower MJ, Aslan CE, Rejmánek M (2009) Invasion potential of *T. sebifera* tree (*Triadica sebifera*) in California's Central Valley. *Invasive Plant Science and Management* 2: 386–395.
- Brisbane City Council (2018) Chinese tallow tree. <https://weeds.brisbane.qld.gov.au/weeds/chinese-tallow-tree>
- Bruce, K. A., Cameron, G. N., & Harcombe, P. A. (1995). Initiation of a new woodland type on the Texas coastal prairie by the Chinese tallow tree (*Sapium sebiferum* (L.) Roxb.). *Bulletin of the Torrey Botanical Club*, 215-225.
- Bruce, K. A., Cameron, G. N., Harcombe, P. A., & Jubinsky, G. (1997). Introduction, impact on native habitats, and management of a woody invader, the Chinese tallow tree, *Sapium sebiferum* (L.) Roxb. *Natural Areas Journal*, 255-260.
- CABI (2017). *Sapium sebiferum* Distribution. <http://www.cabi.org/isc/datasheet/48351>
- Camarillo, S. A., Stovall, J. P., & Sunda, C. J. (2015). The impact of Chinese tallow (*Triadica sebifera*) on stand dynamics in bottomland hardwood forests. *Forest Ecology and Management*, 344, 10-19.
- Cameron, G. N., & Spencer, S. R. (1989). Rapid leaf decay and nutrient release in a Chinese tallow forest. *Oecologia*, 80(2), 222-228.
- Chapman, E. L., Chambers, J. Q., Ribbeck, K. F., Baker, D. B., Tobler, M. A., Zeng, H., & White, D. A. (2008). Hurricane Katrina impacts on forest trees of Louisiana's Pearl River basin. *Forest Ecology and Management*, 256(5), 883-889.
- Conway, W. C.; Smith, L. M.; Bergan, J. F. 1995. Chinese tallow control and the use of tallow woodlands and seeds by avifauna on the Texas Gulf Coast. In: Wester, David B.; Britton, Carlton M., eds. *Research highlights--Noxious brush and weed control; range, wildlife and fisheries management*. Volume 26. Lubbock, TX: Texas Tech University, College of Agricultural Sciences and Natural Resources: 18.
- Cotton TB, Kwiatkowski MA, Saenz D, Collyer M (2012). Effects of an invasive plant, Chinese tallow (*Triadica sebifera*), on development and survival of anuran larvae. *Journal of Herpetology*, 46, 186–193, 201.

DeWalt SJ, Siemann E, Rogers WE (2011). Geographic distribution of genetic variation among native and introduced populations of Chinese tallow tree, *Triadica sebifera* (Euphorbiaceae). *American Journal of Botany*, 98(7), 1128-1138.

EddMapps (2017). Chinese Tallow Distribution.

<https://www.eddmapps.org/distribution/usstate.cfm?sub=3079>

Dillwyn LW, Collinson P (1843) *Hortus Collinsonianus: An account of the plants cultivated by the late Peter Collinson. Arranged alphabetically according to their modern names, from the catalogue of his garden, and other manuscripts, 64 pp.* - Publisher Printed by W. C. Murray and D. Rees. Book digitized by Google from the library of Oxford University and uploaded to the Internet Archive by user tpb. - Source:

<http://books.google.com/books?id=gEQAAAAAQAAJ&oe=UTF-8>

EPPO (2009) EPPO guidelines on the development of a code of conduct on horticulture and invasive alien plants. *EPPO Bulletin* 39, 263–266.

EPPO (2006) PM3/67 Guidelines for the management of invasive alien plants or potentially invasive alien plants which are intended for import or have been intentionally imported. *EPPO Bulletin*, 36, 417-418.

Esser HJ, van Welzen P & Djarwaningsih T (1997) A phylogenetic classification of the Malesian Hippomaneae (Euphorbiaceae). *Systematic Botany* **22**, 617–628

FLEPPC (2005) List of Florida's Invasive Species. Florida Exotic Pest Plant Council. . Accessed 17 Mar 2017.

Flora of China (1972) <http://www.efloras.org/index.aspx>

Gan J, Miller JH, Wang H, Taylor JW (2009). Invasion of tallow tree into southern US forests: influencing factors and implications for mitigation. *Canadian journal of forest research*, 39(7), 1346-1356.

GBIF (2017). *Triadica sebifera* Distribution. <https://www.gbif.org/species/3054399>

Gao R, Su Z, Yin Y and Li S (2016). Germplasm, chemical constituents, biological activities, utilization, and control of Chinese tallow (*Triadica sebifera* (L.) Small). *Biol Invasions* 18:809–829

Grace JB (1998). Can prescribed fire save the endangered coastal prairie ecosystem from Chinese tallow invasion. *Endangered Species Update*, 15(5), 70-76.

Grace JB, Smith MD, Grace SL, Collins SL, Stohlgren TJ (2000). Interactions between fire and invasive plants in temperate grasslands of North America. In *Proceedings of the invasive species workshop: the role of fire in the control and spread of invasive species. Fire conference* (pp. 40-65).

Harcombe PA, Cameron GN, Glumac EG (1993). Above-ground net primary productivity in adjacent grassland and woodland on the coastal prairie of Texas, USA. *Journal of Vegetation Science*, 521-530.

Hortus collinsonianus (1843) an Account of the Plants cultivated by the late Peter Collinson, Esq., F.R.S. available at:

https://books.google.fr/books?id=gEQAAAAAQAAJ&pg=PR9&source=gbs_selected_pages&ad=2#v=onepage&q&f=false

Hortus Paddingtonensis (1797) A catalogue of plants cultivated in the garden of J. Symmons, Esq., Paddington-House. Available at: <https://archive.org/details/b22386890>

Jaryan V, Uniyal SK, Kumar A, Gupta RC, Singh RD (2013). Extent of occurrence and area of occupancy of tallow tree (*Sapium sebiferum*): using the red list criteria for documenting invasive species expanse. National Academy Science Letters, 36(1), 85-91.

Jones RH, McLeod KW (1989). Shade tolerance in seedlings of Chinese tallow tree, American sycamore, and cherrybark oak. Bulletin of the Torrey Botanical Club, 371-377.

Jubinsky G, Anderson LC (1996). The invasive potential of Chinese tallow-tree (*Sapium sebiferum* Roxb.) in the Southeast. Castanea, 226-231.

Kirmse RD, Fisher JT (1989) Species screening and biomass trials of woody plants in the semi-arid southwest United States. Biomass. 18(1): 15-29.

Langeland KA (2009) Natural area weeds: Chinese tallow (*Sapium sebiferum* L.), [Online]. In: EDIS (Electronic Data Information Source). Publication #SS-AGR-45. Gainesville, FL: University of Florida, IFAS (Institute of Food and Agricultural Science) Extension (Producer). Available: <http://edis.ifas.ufl.edu/ag148>

Lieux MH (1975). Dominant pollen types recovered from commercial Louisiana honeys. Economic Botany, 29(1), 87-96.

Leonard, N. (2008) The effects of the invasive exotic Chinese tallow tree (*Triadica sebifera*) on amphibians and aquatic invertebrates. PhD Thesis, University of New Orleans, USA.

Lin WC, Chen AC, Tseng CJ, Hwang SG (1958). An investigation and study of Chinese tallow tree in Taiwan (*Sapium sebiferum*, Roxb.). Bulletin of Taiwan Forestry Research Institute, 57, 32-37.

Loiseleur-Deslongchamps J (1832) Rapport sur les cultures qu'il serait utiles d'introduire ou de perfectionner dans la colonie d'Alger. Book digitized by Google, Source: https://books.google.it/books?id=6g7l-08hejkC&hl=it&source=gbp_navlinks_s

Low T, Booth C (2007) The weedy truth about biofuels. Invasive Species Council, Melbourne, Australia.

McCormick, C. (2005). Chinese Tallow Management Plan for Florida: Recommendations from The Chinese Tallow Task Force. The Florida Exotic Pest Plant Council.

Meyer R (2011) *Triadica sebifera*. In: Fire Effects Information System, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: <http://www.fs.fed.us/database/feis/> [2017, August 30]

Neyland R, Meyer HA (1997). Species diversity of Louisiana chenier woody vegetation remnants. Journal of the Torrey Botanical Society, 254-261.

Nijjer S, Lankau RA, Rogers WE, Siemann E (2002). Effects of temperature and light on Chinese tallow (*Sapium sebiferum*) and Texas sugarberry (*Celtis laevigata*) seed germination. Texas Journal of Science, 54(1), 63-68.

NSW Government, Australia (2017). NSW WeedWise: Chinese tallow tree (*Triadica sebifera*). Department of Primary Industries. Factsheet: <http://weeds.dpi.nsw.gov.au/Weeds/Details/38>

Pattison RR, Mack RN (2008). Potential distribution of the invasive tree *Triadica sebifera* (Euphorbiaceae) in the United States: evaluating CLIMEX predictions with field trials. *Global Change Biology*, 14(4), 813-826.

Pile LS, Wang GG, Knapp BO, Walker JL, Stambaugh MC (2017) Chinese tallow (*Triadica sebifera*) invasion in maritime forests: the role of anthropogenic disturbance and its management implication.

Pogue MG (2014). A new species of *Gadirtha* Walker (Nolidae, Eligminae): a proposed biological control agent of Chinese tallow (*Triadica sebifera* (L.) Small) (Euphorbiaceae) in the United States. *ZooKeys*, (382), 13.

Radford AE, Ahles HE, Bell CR (1968) *Manual of the vascular flora of the Carolinas*. Chapel Hill, NC: The University of North Carolina Press. Pg. 667

Renne IJ, Barrow WC, Randall J, Lori A, Bridges WC (2002). Generalized avian dispersal syndrome contributes to Chinese tallow tree (*Sapium sebiferum*, Euphorbiaceae) invasiveness. *Diversity and Distributions*, 8(5), 285-295.

Renne, IJ, Gauthreaux Jr, SA, Gresham CA (2000). Seed dispersal of the Chinese tallow tree (*Sapium sebiferum* (L.) Roxb.) by birds in coastal South Carolina. *The American Midland Naturalist*, 144(1), 202-215.

Renne IJ, Spira TP, Bridges Jr, WC. (2001). Effects of habitat, burial, age and passage through birds on germination and establishment of Chinese tallow tree in coastal South Carolina. *Journal of the Torrey Botanical Society*, 109-119.

Rogers WE, Siemann E (2002). Effects of simulated herbivory and resource availability on native and invasive exotic tree seedlings. *Basic and Applied Ecology*, 3(4), 297-307.

Rogers, WE, Siemann E (2005). Herbivory tolerance and compensatory differences in native and invasive ecotypes of Chinese tallow tree (*Sapium sebiferum*). *Plant Ecology*, 181(1), 57-68.

Roy, H.E., Adriaens, T., Aldridge, D.C., Bacher, S., Bishop, J.D.D., Blackburn, T.M., Branquart, E., Brodie, J., Carboneras, C., Cook, E.J., Copp, G.H., Dean, H.J., Eilenberg, J., Essl, F., Gallardo, B., Garcia, M., García-Berthou, E., Genovesi, P., Hulme, P.E., Kenis, M., Kerckhof, F., Kettunen, M., Minchin, D., Nentwig, W., Nieto, A., Pergl, J., Pescott, O., Peyton, J., Preda, C., Rabitsch, W., Roques, A., Rorke, S., Scalera, R., Schindler, S., Schönrogge, K., Sewell, J., Solarz, W., Stewart, A., Tricarico, E., Vanderhoeven, S., van der Velde, G., Vilà, M., Wood, C.A., Zenetos, A. (2015) *Invasive Alien Species - Prioritising prevention efforts through horizon scanning ENV.B.2/ETU/2014/0016*. European Commission.

Scheld HW, Cowles JR (1981). Woody biomass potential of the Chinese tallow tree. *Economic Botany*, 35(4), 391-397.

Sharma S, Rikhari HC, PALNI LS (1996). Adoption of a potential plantation tree crop as an agroforestry species but for the wrong reasons: a case study of the Chinese tallow tree from central Himalaya. *International Tree Crops Journal*, 9(1), 37-45.

Siemann, E, Rogers WE (2001) Genetic differences in growth of an invasive tree

species. *Ecology Letters*. 4(6): 514-518. [54558]

Siemann, E, Rogers WE (2003a) Herbivory, disease, recruitment limitation, and success of alien and native tree species. *Ecology*. 84(6): 1489-1505. [45140]

Siemann E, Rogers, WE (2003b) Increased competitive ability of an invasive tree may be limited by an invasive beetle. *Ecological Applications*. 13(6): 1503-1507. [54559]

Siemann, E; Rogers WE (2003c) Reduced resistance of invasive varieties of the alien tree *Sapium sebiferum* to a generalist herbivore. *Oecologia*. 135(3): 451-457. [54560]

Siemann, E; Rogers, WE. Grace JB (2007) Effects of nutrient loading and extreme rainfall events on coastal tallgrass prairies: invasion intensity, vegetation responses, and carbon and nitrogen distribution. *Global Change Biology*. 13: 2184-2192. [79450]

Siemann, E, Carrillo JA, Gabler CA, Zipp R, Rogers WE (2009). Experimental test of the impacts of feral hogs on forest dynamics and processes in the southeastern US. *Forest ecology and management*, 258(5), 546-553.

Société nationale d'horticulture de France (1832) Jardins de France, Volumes 10-11, Book digitized by Google, Source: <https://books.google.it/books?id=xh9OAAAAYAAJ>

USDA (2008). Plant guide: Chinese tallow tree--*Triadica sebifera* (L.) Small, [Online]. In: PLANTS Database--fact sheets & plant guides. Baton Rouge, LA: U.S. Department of Agriculture, Natural Resources Conservation Service, National Plant Data Center (Producer). Available: http://plants.usda.gov/plantguide/pdf/pg_trse6.pdf

Vossen, H.A.M. van der and B.E. Umali (ed)(2001), , Plant Resources of South-East Asia 14. Vegetable oils and fats, Backhuys Publishers, Leiden, The Netherlands, 229 pp

Wang H. H., Grant, W. E., Swannack, T. M., Gan, J., Rogers, W. E., Koralewski, T. E., ... & Taylor, J. W. (2011). Predicted range expansion of Chinese tallow tree (*Triadica sebifera*) in forestlands of the southern United States. *Diversity and Distributions*, 17(3), 552-565.

Wen L, Wang, Y, Lu D, Hu S, Han H (2010). Preparation of KF/CaO nanocatalyst and its application in biodiesel production from Chinese tallow seed oil. *Fuel*, 89(9), 2267-2271.

Wheeler GS, Ding J (2014). Is Chinese tallowtree, *Triadica sebifera*, an appropriate target for biological control in the United States?

Zou, J. Robers WE. Siemann E (2007) Differences in morphological and physiological traits between native and invasive populations of *Sapium sebiferum*. *Functional Ecology*. 21: 721-730

Zou, Jianwen; Rogers, William E.; DeWalt, Saara J.; Siemann, Evan. 2006. The effect of Chinese tallow tree (*Sapium sebiferum*) ecotype on soil-plant system carbon and nitrogen processes. *Oecologia*. 150(2): 272-281.

Zou, J.; Rogers, W. E.; Siemann, E.. 2008. Increased competitive ability and herbivory tolerance in the invasive plant *Sapium sebiferum*. *Biological Invasions*. 10: 291-302.

Zou, J.; Rogers, W. E.; Siemann, E.. 2009. Plasticity of *Sapium sebiferum* seedling growth to light and water resources: inter- and intraspecific comparisons. *Basic and Applied Ecology*. 10: 79-88.

Appendix 1: Projection of climatic suitability for *Triadica sebifera* establishment

Aim

To project the suitability for potential establishment of *Triadica sebifera* in the EPPO region, under current and predicted future climatic conditions.

Data for modelling

Climate data were taken from ‘Bioclim’ variables contained within the WorldClim database (Hijmans *et al.* 2005) originally at 5 arcminute resolution (0.083 x 0.083 degrees of longitude/latitude) and aggregated to a 0.25 x 0.25 degree grid for use in the model. Based on the biology of the focal species, the following climate variables were used in the modelling:

- Mean minimum temperature of the coldest month (Bio6 °C) reflecting exposure to frost. *Triadica sebifera* suffers substantial damage after exposure to freezing temperatures for 36 hours or more (Grace *et al.* 2000). In its invaded range in North America, *T. sebifera* did not occur at survey sites where the mean minimum temperature in January was below – 12 °C (Gan *et al.* 2009).
- Mean temperature of the warmest quarter (Bio10 °C) reflecting the growing season thermal regime. Previous CLIMEX modelling inferred a lower temperature threshold for growth of 12 °C, optimal growing conditions between 24 and 35 °C and an upper temperature threshold of 40 °C (Pattison & Mack, 2008).
- Climatic moisture index (CMI, ratio of mean annual precipitation, Bio12, to PET) reflecting plant moisture regimes. *Triadica sebifera* occurrence may be restricted by drought since plantations in arid regions are irrigated (Kirmse & Fisher, 1989).
- Temperature seasonality (Bio15, coefficient of variation for monthly precipitations) as another measure of the moisture regime. *Triadica sebifera* is likely to favour even precipitation regimes where periods of drought stress are minimal.
- Precipitation of the coldest quarter (Bio19, mm) as winter recharge of soil moisture is likely to minimise drought stress of germinating seedlings in spring.

To estimate the effect of climate change on the potential distribution, equivalent modelled future climate conditions for the 2070s under the Representative Concentration Pathway (RCP) 4.5 and 8.5 were also obtained. For both scenarios, the above variables were obtained as averages of outputs of eight Global Climate Models (BCC-CSM1-1, CCSM4, GISS-E2-R, HadGEM2-AO, IPSL-CM5A-LR, MIROC-ESM, MRI-CGCM3, NorESM1-M), downscaled and calibrated against the WorldClim baseline (see http://www.worldclim.org/cmip5_5m).

RCP 4.5 is a moderate climate change scenario in which CO₂ concentrations increase to approximately 575 ppm by the 2070s and then stabilise, resulting in a modelled global temperature rise of 1.8 C by 2100. RCP8.5 is the most extreme of the RCP scenarios, and may therefore represent the worst case scenario for reasonably anticipated climate change. In RCP8.5 atmospheric CO₂ concentrations increase to approximately 850 ppm by the 2070s, resulting in a modelled global mean temperature rise of 3.7 °C by 2100.

In the models the following habitat variables were also included:

- Human influence index as *T. sebifera*, like many invasive species, is likely to associate with anthropogenically disturbed habitats. We used the Global Human Influence Index Dataset of the Last of the Wild Project (Wildlife Conservation Society - WCS & Center for International Earth Science Information Network - CIESIN - Columbia University, 2005), which is developed from nine global data layers covering human population pressure (population density), human land use and infrastructure (built-up areas, nighttime lights,

land use/land cover) and human access (coastlines, roads, railroads, navigable rivers). The index ranges between 0 and 1 and was log+1 transformed for the modelling to improve normality.

- Tree cover as *T. sebifera* may be restricted from closed canopy forest (Bruce, 1993, Jones & McLeod, 1989). Tree cover was estimated from the MODerate-resolution Imaging Spectroradiometer (MODIS) satellite continuous tree cover raster product, produced by the Global Land Cover Facility (<http://glcf.umd.edu/data/vcf/>). The raw product contains the percentage cover by trees in each 0.002083 x 0.002083 degree grid cell, from which mean cover was aggregated to the 0.25 x 0.25 degree grid.
- Density of permanent rivers was estimated from the Vector Map (VMAPO; <http://gis-lab.info/qa/vmap0-eng.html>). River vectors were rasterised at 0.02 x 0.02 degree resolution. Then, the percentage of these grid cells containing rivers within each of the 0.25 x 0.25 degree cells used in the model was calculated.

Species occurrence data were obtained from the Global Biodiversity Information Facility (GBIF), USDA Biodiversity Information Serving Our Nation (BISON), Berkeley Ecoinformatics Engine and Eddmaps. Occurrence records were scrutinised to remove those from regions where the species is not known to be well established, those that appeared to be dubious or planted specimens (e.g. plantations, botanic gardens) and those where the georeferencing was too imprecise (e.g. records referenced to a country or island centroid) or outside of the coverage of the predictor layers (e.g. small island or coastal occurrences). The remaining records were gridded at a 0.25 x 0.25 degree resolution for modelling (Figure 1a). In total 858 grid cells contained records of *T. sebifera*.

Additionally, the recording density of vascular plants (phylum Tracheophyta) on GBIF was obtained as a proxy for spatial recording effort bias (Figure 1b).




Figure 1. (a) Occurrence records obtained for *Triadica sebifera* and used in the modelling, showing the native range and (b) a proxy for recording effort – the number of Tracheophyta records held by the Global Biodiversity Information Facility, displayed on a \log_{10} scale.

Species distribution model

A presence-background (presence-only) ensemble modelling strategy was employed using the BIOMOD2 R package v3.3-7 (Thuiller *et al.* 2014, Thuiller *et al.* 2009). These models contrast the environment at the species' occurrence locations against a random sample of background environmental conditions (often termed 'pseudo-absences') in order to characterise and project suitability for occurrence. This approach has been developed for distributions that are in equilibrium with the environment. Because invasive species' distributions are not at equilibrium and subject to dispersal constraints at a global scale, we took care to minimise the inclusion of locations suitable for the species but where it has not been able to disperse to. Therefore the background sampling region included:

- The area accessible by native *T. sebifera* populations (see Fig. 1a), in which the species is likely to have had sufficient time to disperse to all locations. The accessible native region was defined as a 300 km buffer around the minimum convex polygon bounding all native occurrences in East Asia; AND
- A relatively small 30 km buffer around all non-native occurrences, encompassing regions likely to have had high propagule pressure for introduction by humans and/or dispersal of the species; AND
- Regions where we have an *a priori* expectation of high unsuitability for the species (see Figure 2). Absence from these regions is considered to be irrespective of dispersal constraints. Based on published ecophysiological information described above and the extremes of the climatic predictors at the species occurrences the following rules for unsuitability were applied:
 - Mean minimum temperature of the coldest month (Bio6) < -5 °C.
 - Mean temperature of the warmest quarter (Bio10) < 16 °C.
 - Mean temperature of the warmest quarter (Bio10) > 29 °C.
 - Climatic moisture index (CMI) < 0.45.
 - Precipitation of coldest quarter (Bio19) < 20 mm.

In all, 3.7 % of occurrence grid cells exceeded any one of these thresholds. From this background region, ten samples of 10,000 randomly chosen grid cells were obtained (Figure 2). To account for recording effort bias, sampling of background grid cells was weighted in proportion to the Tracheophyte recording density (Figure 1b).




Figure 2. Randomly selected background grid cells used in the modelling of *Triadica sebifera*, mapped as black points. Points are sampled from the native range, a small buffer around non-native occurrences and from areas expected to be highly unsuitable for the species (grey background region), and weighted by a proxy for plant recording effort (Figure 1b).

Each dataset (i.e. combination of the presences and the individual background samples) was randomly split into 80% for model training and 20% for model evaluation. With each training dataset, ten statistical algorithms were fitted with the default BIOMOD2 settings (except where specified below) and rescaled using logistic regression:

- Generalised linear model (GLM)
- Generalised boosting model (GBM)
- Generalised additive model (GAM) with a maximum of four degrees of freedom per effect.
- Classification tree algorithm (CTA)
- Artificial neural network (ANN)
- Flexible discriminant analysis (FDA)
- Multivariate adaptive regression splines (MARS)
- Random forest (RF)
- MaxEnt
- Maximum entropy multinomial logistic regression (MEMLR)

Since the background sample was much larger than the number of occurrences, prevalence fitting weights were applied to give equal overall importance to the occurrences and the background. Normalised variable importance was assessed and variable response functions were produced using BIOMOD2's default procedure. Model predictive performance was assessed by calculating the Area Under the Receiver-Operator Curve (AUC) for model predictions on the evaluation data, that were reserved from model fitting. AUC can be interpreted as the probability that a randomly selected presence has a higher model-predicted suitability than a randomly selected absence.

An ensemble model was created by first rejecting poorly performing algorithms with relatively extreme low AUC values and then averaging the predictions of the remaining algorithms, weighted by their AUC. To identify poorly performing algorithms, AUC values were converted into modified z-scores based on their difference to the median and the median absolute deviation across all algorithms (Iglewicz & Hoaglin, 1993). Algorithms with $z < -2$ were

rejected. In this way, ensemble projections were made for each dataset and then averaged to give an overall suitability.

Global model projections were made for the current climate and for the two climate change scenarios, avoiding model extrapolation beyond the ranges of the input variables. The optimal threshold for partitioning the ensemble predictions into suitable and unsuitable regions was determined using the ‘minimum ROC distance’ method. This finds the threshold where the Receiver-Operator Curve (ROC) is closest to its top left corner, i.e. the point where the false positive rate (one minus specificity) is zero and true positive rate (sensitivity) is one.

Limiting factor maps were produced following Elith *et al.* (2010). For this, projections were made separately with each individual variable fixed at a near-optimal value. These were chosen as the median values at the occurrence grid cells. Then, the most strongly limiting factors were identified as the one resulting in the highest increase in suitability in each grid cell. Partial response plots were also produced by predicting suitability across the range of each predictor, with other variables held at near-optimal values.

Results

The ensemble model suggested that suitability for *T. sebifera* was most strongly determined by summer temperature, moisture availability and winter temperature (Table 1). From Figure 3, suitability was restricted by low summer temperature, drought and low winter temperatures. A weaker preference for human-influenced regions, wet winters, and even precipitation regimes was also modelled, but tree cover and river density had very little effect at the scale of the model. For all predictors, there was substantial variation in the partial response plots between algorithms (Figure 3).

Global projection of the model in current climatic conditions indicates that the main clusters of native and known invaded records fell within regions predicted to have high suitability (Figure 4). The model predicts potential for further northwards expansion of the non-native range in North America, as well as in South America and southern Africa, where there are currently very few records of invasive populations (Figure 4).

The model predicts a region of potential suitability for *T. sebifera* in southern Europe (Figure 5). Countries in which *T. sebifera* may be capable of establishing widely include all those bordering the northern Mediterranean Sea, from Portugal to Turkey, as well as the Black Sea coasts of Turkey, Georgia and Russia. The model predicts that establishment in the rest of Europe will largely be prevented because of low summer temperature, with moisture limitation in central Spain and frost limitation in far eastern central Europe (Figure 6).

The climate change projections for the 2070s suggest that under the less extreme RCP4.5 scenario the suitable region in Europe will have extended northwards, especially in western Europe where warmer summers mean the species may be capable of establishing in southern United Kingdom and the southern Baltic coast (Figure 7). Under the more extreme RCP8.5 scenario, the northwards expansion is even greater, especially in eastern Europe (Figure 8). This is likely driven by a relaxation of frost constraints.

In terms of Biogeographical Regions, those predicted to be most suitable for *T. sebifera* establishment in the current climate are the Mediterranean and Black Sea (Figure 9). In the evaluated climate change scenarios, predicted suitability was stable in the Mediterranean but increased in the Black Sea. Other biogeographic regions predicted to strongly increase in suitability are Atlantic, Continental, Pannonian and Steppic (Figure 9).

Table 1. Summary of the cross-validation predictive performance (AUC) and variable importances of the fitted model algorithms and the ensemble (AUC-weighted average of the best performing algorithms). Results are the average from models fitted to ten different background samples of the data.

Algorithm	Predictive AUC	In the ensemble	Variable importance						
			Minimum temperature of coldest month	Mean temperature of warmest quarter	Climatic moisture index	Precipitation seasonality	Precipitation of coldest quarter	Human influence index	Tree cover
GBM	0.9797	yes	13%	47%	32%	0%	6%	1%	0%
GAM	0.9773	yes	18%	42%	24%	1%	3%	11%	1%
ANN	0.9765	yes	21%	38%	26%	2%	4%	6%	4%
MARS	0.9752	yes	17%	44%	29%	1%	3%	5%	0%
Maxent	0.9713	yes	17%	45%	17%	7%	8%	5%	1%
RF	0.9707	yes	6%	67%	10%	5%	4%	5%	2%
GLM	0.9706	yes	20%	47%	22%	0%	1%	7%	2%
FDA	0.9649	yes	9%	67%	17%	0%	6%	0%	0%
CTA	0.9435	no	19%	38%	30%	4%	7%	2%	1%
MEMLR	0.8507	no	15%	21%	7%	31%	16%	1%	9%
Ensemble	0.9788		15%	50%	22%	2%	5%	5%	1%




Figure 3. Partial response plots from the fitted models, ordered from most to least important. Thin coloured lines show responses from the algorithms in the ensemble, while the thick black line is their ensemble. In each plot, other model variables are held at their median value in the training data. Some of the divergence among algorithms is because of their different treatment of interactions among variables.




Figure 4. (a) Projected global suitability for *Triadica sebifera* establishment in the current climate. For visualisation, the projection has been aggregated to a 0.5 x 0.5 degree resolution, by taking the maximum suitability of constituent higher resolution grid cells. Red shading indicates suitability. White areas have climatic conditions outside the range of the training data so were excluded from the projection. (b) Uncertainty in the suitability projections, expressed as the standard deviation of projections from different algorithms in the ensemble model.




Figure 5. Projected current suitability for *Triadica sebifera* establishment in Europe and the Mediterranean region. The white areas have climatic conditions outside the range of the training data so were excluded from the projection.




Figure 6. Limiting factor map for *Triadica sebifera* establishment in Europe and the Mediterranean region in the current climate. Shading shows the predictor variable most strongly limiting projected suitability, which is most informative for places predicted unsuitable for the species. See section ‘data for modelling’ for information relating to Bio6-19.




Figure 7. Projected suitability for *Triadica sebifera* establishment in Europe and the Mediterranean region in the 2070s under climate change scenario RCP4.5, equivalent to Figure 5.




Figure 8. Projected suitability for *Triadica sebifera* establishment in Europe and the Mediterranean region in the 2070s under climate change scenario RCP8.5, equivalent to Figure 5.





Figure 9. Variation in projected suitability among Biogeographical regions of Europe. The bar plots show the proportion of grid cells in each region classified as suitable in the current climate and projected climate for the 2070s under emissions scenarios RCP4.5 and RCP8.5. The coverage of each region is shown in the map below.

Caveats to the modelling

Modelling the potential distributions of range-expanding species is always difficult and uncertain. We did not have sufficient information to determine whether all records used in the modelling were from established populations rather than plantations. If the latter were included, management to alleviate climatic stresses (such as irrigation) may have caused the model to over-estimate the niche breadth and potential establishment distribution of *T. sebifera*.

Triadica sebifera is largely restricted to wet micro-habitats such as river banks in the more arid parts of its global distribution (e.g. California and Mexico), so it is likely that establishment in the more arid parts of the suitable region in Europe would also be restricted to such habitats. Although we attempted to model this interaction by including river density in the model, local habitat factors are unlikely to be well represented at the scale of the model.

The limiting factors map may have under-estimated the limiting influence of winter temperatures in Europe, since two of the algorithms in the ensemble did not model a strong limitation of suitability at very cold temperatures. This will have raising the ensemble model suitability response to very cold winter temperatures.

Other variables potentially affecting the distribution of the species, such as edaphic variables, were not included in the model.

To remove spatial recording biases, the selection of the background sample was weighted by the density of Tracheophyte records on the Global Biodiversity Information Facility (GBIF). While this is preferable to not accounting for recording bias at all, a number of factors mean this may not be the perfect null model for species occurrence:

- The GBIF API query used to did not appear to give completely accurate results. For example, in a small number of cases, GBIF indicated no Tracheophyte records in grid cells in which it also yielded records of the focal species.
- Additional data sources to GBIF were used, which may have been from regions without GBIF records.

References

Hijmans RJ, Cameron SE, Parra JL, Jones PJ, Jarvis A (2005) Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology* **25**, 1965-1978.

Grace JB, Smith MD, Grace SL, Collins SL, Stohlgren TJ (2000) Interactions between fire and invasive plants in temperate grasslands of North America. In *Proceedings of the invasive species workshop: the role of fire in the control and spread of invasive species. Fire conference*, pp. 40-65.

Gan J, Miller JH, Wang H, Taylor JW (2009) Invasion of tallow tree into southern US forests: influencing factors and implications for mitigation. *Canadian journal of forest research* **39**, 1346-1356.

Pattison RR, Mack RN (2008) Potential distribution of the invasive tree *Triadica sebifera* (Euphorbiaceae) in the United States: evaluating CLIMEX predictions with field trials. *Global Change Biology* **14**, 813-826.

Kirmse RD, Fisher JT (1989) Species screening and biomass trials of woody plants in the semi-arid southwest United States. *Biomass* **18**, 15-29.

Wildlife Conservation Society - WCS & Center for International Earth Science Information Network - CIESIN - Columbia University (2005) Last of the Wild Project, Version 2, 2005 (LWP-2): Global Human Influence Index (HII) Dataset (Geographic). NASA Socioeconomic Data and Applications Center (SEDAC), Palisades, NY.

Bruce KA (1993) Factors affecting the biological invasion of the exotic Chinese tallow tree, *Sapium sebiferum*, in the Gulf Coast Prarie of Texas. University of Houston.


Jones RH, McLeod KW (1989) Shade tolerance in seedlings of Chinese tallow tree, American sycamore, and cherrybark oak. *Bulletin of the Torrey Botanical Club*, 371-377.

Thuiller W, Georges D, Engler R (2014) biomod2: Ensemble platform for species distribution modeling. R package version 3.3-7 Available at: <https://cran.r-project.org/web/packages/biomod2/index.html>.

Thuiller W, Lafourcade B, Engler R, Araújo MB (2009) BIOMOD—a platform for ensemble forecasting of species distributions. *Ecography* **32**, 369-373.

Iglewicz B, Hoaglin DC (1993) *How to detect and handle outliers*. Asq Press.

Appendix 2. Biogeographical regions



Appendix 3 Images



Figure 1. *Triadica sebifera* invasion in North America



Figure 2. *Triadica sebifera* trunk



Figure 3. *Triadica sebifera* fruits

Appendix 4: Distribution summary for EU Member States and Biogeographical regions

Member States:

	Recorded	Established (currently)	Established (future)	Invasive (currently)
Austria	–	–	–	–
Belgium	–	–	–	–
Bulgaria	–	–	–	–
Croatia	–	–	YES	–
Cyprus	–	–	–	–
Czech Republic	–	–	–	–
Denmark	–	–	–	–
Estonia	–	–	–	–
Finland	–	–	–	–
France	–	–	YES	–
Germany	–	–	–	–
Greece	–	–	YES	–
Hungary	–	–	–	–
Ireland	–	–	–	–
Italy	–	–	YES	–
Latvia	–	–	–	–
Lithuania	–	–	–	–
Luxembourg	–	–	–	–
Malta	–	–	–	–
Netherlands	–	–	–	–
Poland	–	–	–	–
Portugal	–	–	YES	–
Romania	–	–	–	–
Slovakia	–	–	–	–
Slovenia	–	–	–	–
Spain	–	–	YES	–
Sweden	–	–	–	–
United Kingdom	–	–	–	–

Biogeographical regions

	Recorded	Established (currently)	Established (future)	Invasive (currently)
Alpine	–	–	–	–
Atlantic	–	–	YES	–
Black Sea	–	–	–	–
Boreal	–	–	–	–
Continental	–	–	YES	–
Mediterranean	–	–	YES	–
Pannonian	–	–	–	–
Steppic	–	–	–	–

YES: if recorded in natural environment, established or invasive or can occur under future climate; – if not recorded, established or invasive; ? Unknown

Appendix 5 Maps⁴




Figure 1. Global distribution of *Triadica sebifera*

⁴ Note Maps in appendix 5 may contain records, e.g. herbarium records, that were not considered during the climate modelling stage. Data sources are from literature, GBIF and expert opinion.




Figure 2. North American distribution for *Triadica sebifera*




Figure 3. Asia distribution for *Triadica sebifera*




Figure 4. Australia distribution for *Triadica sebifera*